

# Rhodora

JOURNAL OF THE  
NEW ENGLAND BOTANICAL CLUB.

Conducted and published for the Club, by

BENJAMIN LINCOLN ROBINSON Editor-in-chief.

FRANK SHIPLEY COLLINS

MERRITT LYNDON FERNALD } Associate Editors.

HOLLIS WEBSTER

WILLIAM PENN RICH

EDWARD LOTHROP RAND } Publication Committee.

Vol. 13.

December, 1911.

No. 156.

## CONTENTS:

Typhae of Maryland and Virginia. <i>Ivar Tidestrom</i> . . . . .	241
Two Lost Carices of eastern Massachusetts. <i>M. L. Fernald</i> . . . . .	243
Flora of the Boston District,—XIII . . . . .	248
<i>Salix calcicola</i> . <i>M. L. Fernald and K. M. Wiegand</i> . . . . .	251
Stump-healing in <i>Pinus Strobus</i> . <i>A. J. Eames</i> . . . . .	253
Botanical Lectures at the Lowell Institute . . . . .	253
Errata . . . . .	254
Index . . . . .	255

Boston, Mass.  
1052 Exchange Building.



Providence, R. I.  
Preston and Rounds Co.

**RHODORA**.—A monthly journal of botany, devoted primarily to the flora of New England. Price \$1.00 per year (\$1.25 to all foreign countries including Canada); single copies 15 cents. Volume 1, \$2.00, Vol. 2, \$1.50. All remittances by check or draft, except on Boston or New York, must include ten cents additional for cost of collection. Notes and short scientific papers, relating directly or indirectly to the plants of the northeastern states, will be gladly received and published to the extent that the limited space of the journal permits. Forms will be closed five weeks in advance of publication. Authors (of more than one page of print) will receive 25 copies of the issue in which their contributions appear. Extracted reprints, if ordered in advance, will be furnished at cost.

Address manuscripts and proofs to

B. L. ROBINSON, 3 Clement Circle, Cambridge, Mass.

Subscriptions, advertisements, and business communications to

W. P. RICH, 300 Massachusetts Avenue, Boston, Mass.

Single copies may be had from

E. L. RAND, Corresponding Sec'y N. E. Botanical Club,  
1052 Exchange Building, Boston, Mass.

Entered at Boston, Mass., Post office as Second Class Mail Matter

---

**KEY TO NEW ENGLAND TREES**, Wild and Commonly Cultivated, based primarily upon leaf characters, by J. FRANKLIN COLLINS and HOWARD W. PRESTON. Price 40c. net. PRESTON & ROUNDS CO., Providence, R. I.

---

**CARD-INDEX OF NEW GENERA, SPECIES AND VARIETIES OF AMERICAN PLANTS, 1885 TO DATE.**

For American taxonomists and all students of American plants the most important supplement to the Index Kewensis, this catalogue in several ways exceeds the latter work in detail, since it lists not merely the flowering plants, but pteridophytes and cellular cryptogams, and includes not merely genera and species, but likewise subspecies, varieties and forms. A work of reference invaluable for larger herbaria, leading libraries, academies of sciences, and other centers of botanical activity. Issued quarterly, at \$15.00 per 1000 cards.

GRAY HERBARIUM of Harvard University,  
Cambridge, Mass., U. S. A.

---

**CHECK LIST OF GRAY'S MANUAL, 7th EDITION**, compiled by M. A. DAY. Leatherette. Pocket size. Invaluable for collectors' memoranda and herbarium records. Published and sold by the GRAY HERBARIUM, Cambridge, Mass. Price post-paid 20 cts. each. Ten copies \$1.50.

---

*Advertisements of Nurserymen and Dealers in Botanical and other Scientific Publications are inserted in these pages at the following rates per space of 4 in. by 3-4 in. 1 year \$4.00, 6 months \$2.50.*

# Rhodora

JOURNAL OF

THE NEW ENGLAND BOTANICAL CLUB

---

Vol. 13.

December, 1911.

No. 156.

---

## THE TYPHAE OF MARYLAND AND VIRGINIA.

IVAR TIDESTROM.

In the manuals of recent dates<sup>1</sup> and also in the North American Flora<sup>2</sup> the two circumboreal species *Typha latifolia* and *T. angustifolia* are credited to our region. These are readily distinguished by their floral structures and particularly by the pollen grains — the latter having simple grains while in the former they are in tetrads. Besides, the commonly contiguous spikes in *T. latifolia* and its much broader leaves serve to distinguish it from *T. angustifolia*. In the latter the spikes are separated by an interval of from 1 to 5 cm. or more and the leaves, at least in the normal form, are only one-half as wide or about 6 mm. more or less. There are other characters which need not be mentioned, since they are given in the manuals.

In the *Bracteolatae* or *T. angustifolia* group there are some variants which approach *T. latifolia* in outward appearance but have the floral structure and remote spikes of the other species. Of these we find one in Maryland and Virginia.

On October 17, 1909, Dr. C. L. Alsberg and myself, while botanizing in the neighborhood of Cape Henry, came upon an area of *Typha* about one mile or more west of the lighthouse. At first I thought we had *T. latifolia* but upon closer examination I noticed that we had a plant before us with leaves as broad as those of *T. latifolia* and the remote spikes of the other species. I remarked to my companion that we had a new form or perhaps an old forgotten one before us. Since the variation in the size and form of our water- or marsh-plants ranges between rather wide limits, I did not wish to propose a new

<sup>1</sup> Britton, Man. 38, 1905. Gray's Man. 68, 1908.

<sup>2</sup> N. Am. Fl. 17<sup>1</sup>: 3-4, 1909.



form until more was known about the distribution of the plant and flowering specimens had been collected.

On August 3-4, this year, Professor A. S. Hitchcock, one of his sons and myself botanized between Leonardtown and Millstone, St. Mary's County, Md., and on this trip we located another area of the same form. We were able to refer the plants immediately to the *Bracteolatae*, although they had the appearance of *T. latifolia*. There appear to be no structural characters upon which this plant can be separated from *T. angustifolia*. The broader leaves and somewhat larger spikes can hardly merit more than formal or varietal distinction. If we should accept the views of those whom an eminent French authority has called "les pulverisateurs" or "les mihicists," then a mere difference in the aspect of a plant would compel us to erect a new species. This we cannot do conscientiously in the present case until more of the life history of the plant is known. Rafinesque described a number of species of *Typha*, but these will probably remain indeterminable since he gave no stable characters.

The following summary may serve in the identification of the forms growing in Maryland and Virginia:

*Spikes commonly contiguous.*

*TYPHA LATIFOLIA* L. Sp. Pl. 971, 1753. Kronfeld, Mon. (in Verh. Zool.-bot. Gesellsch. Wien) 176, 1889.

Plants 1.5-2.5 m. in height: leaves flat, 10-20 mm. broad, equalling or exceeding the inflorescence, spikes nearly equal in length (10-30 cm. long); the pistillate dark-brown; pistillate flowers without bracteoles; pollen grains in tetrads.

In marshes and wet places: common.

*Spikes commonly remote.*

*Leaves about 6 mm. wide.*

*TYPHA ANGUSTIFOLIA* L. l. c. Kronfeld, l. c. 150.

Plants 1-1.5 m. in height: leaves plano-convex, exceeding the inflorescence: spikes of nearly equal length (10-30 cm. long), the interval varying from 1 to 5 cm.; the pistillate brown; flowers bracteolate.

In fresh and brackish marshes. I have observed the species in many localities in southern Maryland and on the Eastern Shore of Maryland and Virginia. Mr. McAtee has observed it at Hyattsville, near Washington.

*Leaves about 10 mm. or more in width.*

***Typha angustifolia* var. *virginica*, var. nov.**

Planta robusta 2–2.5 m. alta. Folia glauca planoconvexa ca. 10 mm. lata caulibus multum longiora. Spica feminea cinnamomea 20 mm. lata 20 cm. longa, floribus bracteolata, e spica mascula 1–5 cm. distans. Pollinis granula solitaria.

In brackish marshes: near Cape Henry, Va. (Tm. 3072); with *Myrica cerifera* and *Baccharis halimifolia* in marsh near Millstone (mouth of the Patuxent River), Md. (Tm. 5141) type material.

*Typha glauca* Godr., which was viewed as a hybrid between *T. angustifolia* and *T. latifolia* by Kronfeld, appears to lie between the two species. He characterized it as follows: *Planta robusta, 12–15 dm. alta. Spica masc. et fem. contiguae. Axis spicae masc. pilis linearibus sordido-albis instructus. Pollen. . . . Spica fem. castaneo-brunnea. Flores sine bracteolis; stigmata linearia pilos albos acutos superantia. Fructus. . . . Folia caulium floriferorum laminata, glauca, laminae planae, ad 10 mm. latae, inflorescentiam superantes.*

As pointed out by Kronfeld this form has the habit and ebracteolate flowers of *T. latifolia*, and the color of the spike and form of the stigma of *T. angustifolia*. I have never observed any plant of the above description. Our variety (*virginica*) has nothing in common with *T. latifolia* except the wide leaves. These however are *plano-convex* in cross-section as in typical *T. angustifolia*.

BUREAU OF PLANT INDUSTRY, Washington, D. C.

## TWO LOST CARICES OF EASTERN MASSACHUSETTS.

M. L. FERNALD.

IN 1836, Dewey<sup>1</sup> described from near Boston two plants collected by B. D. Greene and preserved in the Torrey Herbarium, one as *Carex Greeniana* Dewey, the other taken to be *C. binervis* Smith, the latter said to be "probably introduced like *C. panicea*, from Europe; it is very like the *C. binervis* of Europe." Subsequently, it was generally stated that the plants which Dewey described were *C. helodes* Link (*C. laevigata* Smith) and *C. Hornschuchiana* Hoppe (*C. fulva* of Authors, not Good.), though, as indicated below, there has been

<sup>1</sup> Dewey, Am. Jour. Sci., xxx, 61 (1836).



much confusion in the exact interpretation of Dewey's types and consequently in the statements of which one is *C. helodes*, which *C. Hornschuchiana* — a perplexity to which Dewey himself contributed not a little. In 1845, in Wood's *Class Book*,<sup>1</sup> Dewey included both *C. binervis* and *C. Greeniana* with the statement that the latter "Resembles *C. pelva*, Good. [misprint for *C. fulva*], but differs in its fruit and glume." This statement (with *C. pelva* corrected to *C. fulva*) was repeated in subsequent issues of Wood's *Class Book* until 1861, when Dewey reversed the treatment,<sup>2</sup> changing to *C. fulva* Good. with the synonym "*C. binervis* Ed. 1," and *C. laevigata* Smith with the synonym "*C. Greeniana* Ed. 1" and the statement, "This and the last probably introduced from Eur."

Torrey, in his *Monograph of the North American Cyperaceae*,<sup>3</sup> followed Dewey's original publication; but Carey, in Gray's *Manual*,<sup>4</sup> taking his cue possibly from Dewey's statement in the early editions of Wood's *Class Book*, made *C. binervis* Dew., not Smith = *C. laevigata* Smith from "Massachusetts (Tewksbury? *B. D. Greene*): probably introduced"; while *C. Greeniana* Dew. was reduced to *C. fulva* and said to come from a "Pond at Tewksbury."

In the 2nd edition of the *Manual* however, Carey also reversed his treatment<sup>5</sup> and made *C. Greeniana* (not *C. binervis* Dew.) synonymous with *C. laevigata*, and *C. binervis* Dew. synonymous with *C. fulva*; and this understanding of the matter was followed through the three succeeding editions of the *Manual*. In his monumental *Illustrations of the Genus Carex*, Francis Boott<sup>6</sup> treated *C. Greeniana* Dew. as identical with *C. fulva*, and *C. binervis* Dewey as identical with *C. laevigata*; and Bailey, in his *Preliminary Synopsis of North American Carices*,<sup>7</sup> followed Francis Boott and the earlier statements of Dewey in Wood's *Class Book* in treating *C. Greeniana* as *C. fulva*; and in a recent paper upon Newfoundland, Professor Wiegand and the writer,<sup>8</sup> relying upon Boott's *Illustrations* and Bailey's *Synopsis*, spoke of *C. Greeniana* Dewey as probably a form of *C. Hornschuchiana* Hoppe (*C. fulva* Auct.). In discussing the Newfoundland and Anticosti

<sup>1</sup> Dewey in Wood, Cl. Bk. 424 (1845).

<sup>2</sup> Dewey in Wood, Cl. Bk. 764 (1861).

<sup>3</sup> Torr. Ann. Lyc. Nat. Hist. N. Y. iii. 423, 424 (1836).

<sup>4</sup> Carey in Gray, Man. 559 (1848).

<sup>5</sup> Carey, l. c. ed. 2, 528 (1856).

<sup>6</sup> Boott, Ill. Car. iv. 137, 163 (1867).

<sup>7</sup> Bailey, Proc. Am. Acad. xxii. 112 (1886).

<sup>8</sup> Fernald & Wiegand in Fernald, RHODORA, xiii, 130 (1911).

plant which we proposed as a new American variety of *C. Hornschuchiana* (var. *laurentiana* Fernald & Wiegand) we unfortunately overlooked a paper which was published by Mr. K. K. Mackenzie in 1910. In this paper Mr. Mackenzie had given a clear discussion<sup>1</sup> of the same plant (seen by him from Anticosti and Miquelon and identified with the Massachusetts plant of B. D. Greene which had often been called "*C. fulva*") and proposed it as a new species, *C. fulvescens*, with the suggestion that the specimen collected long ago by Greene near Boston was "possibly introduced through wild fowls from further north," and that the old report of *C. fulva* from Newfoundland arose from finding this species there.

In his discussion of *Carex fulvescens* Mackenzie points out that, in spite of the crisscross statements of Dewey, Carey, and others (including the present writer) in regard to the Greene plants, the specimen in the Torrey Herbarium "marked *Carex Greeniana* is a specimen of the European *Carex helodes* Link (*Carex laevigata* Smith) and has the long-acuminate or aristate scales of that species, in this agreeing with Dewey's description, which calls for a plant with cuspidate or mucronate scales." Thus, it may be concluded, the identity of Dewey's *C. Greeniana* is at last definitely and satisfactorily settled.

But the main object of these notes is to direct the attention of local botanists to the two plants, *C. helodes* Link (*C. laevigata* Smith) and *C. Hornschuchiana* Hoppe, var. *laurentiana* Fernald & Wiegand or *C. fulvescens* Mackenzie.<sup>2</sup> There is no evidence that either of these plants has been found in Massachusetts since their discovery by Greene. Whether *Carex helodes* (*C. laevigata*) was indigenous is questionable, since the species has not been found elsewhere on our side of the Atlantic. But the other plant (*C. Hornschuchiana*, var. *laurentiana* or *C. fulvescens*) which Mr. Mackenzie identifies without question with the Newfoundland, Miquelon and Anticosti plant, is strictly American and ordinarily separable by its larger perigynia and sharper scales from the Old World *C. Hornschuchiana*.

As already pointed out by the present writer<sup>3</sup> and as previously surmised by Mackenzie, the plant was known to Goodenough in

<sup>1</sup> Mackenzie, Bull. Torr. Bot. Cl. xxxvii, 239-241 (1910).

<sup>2</sup> Whether the plant is to be considered an American variety of *C. Hornschuchiana*, which seems to Professor Wiegand and the writer the logical treatment of it, or a distinct species, as Mr. Mackenzie interprets it, is a minor question with perfectly sound reasons for either course.

<sup>3</sup> Fernald, RHODORA, xlii. 130 (1911).



1794 "from *America* and *Newfoundland*." The basis for Goodenough's statement was long a mystery and we find the great Francis Boott, just before his death, still impressed with the fact, that "It is remarkable that Goodenough originally received *C. fulva* from America, and that the late Mr. B. D. Greene found it some years ago near Boston, and that no one has since met with it there or elsewhere in the States."<sup>1</sup> The occurrence of the plant in Newfoundland is now clearly demonstrated, and besides from there and the problematic station of B. D. Greene near Boston we know the plant also from bogs or boggy spots on Anticosti and Miquelon, but nowhere on the North American continent. That it has been found elsewhere there can be little question, since Goodenough's report of it from Newfoundland is now verified and he had also seen the plant "from *America*."

In the hope of determining with more exactness the stations where Greene had found his famous specimens the writer appealed to Dr. N. L. Britton, and in reply to these inquiries Mr. Mackenzie has kindly sent a full statement of the data found on the original sheets in the Torrey Herbarium; but unfortunately neither of the sheets indicates anything but "Boston B. D. Greene." Among Greene's own specimens at the Boston Society of Natural History Dr. J. A. Cushman has made a painstaking search and finds only one sheet: nothing to show for *C. helodes*, but a most important sheet of *C. Hornschuchiana*, var. *laurentiana*. This bears a small label through which the specimen is thrust, with the following in different inks but apparently all in Greene's hand: "*C. fulva*. B. D. Greene [followed by a space where a word is now torn out] but not *limosa* growing with it in pond at Tewks<sup>y</sup>?" and in pencil after the last word (hand unidentified): "certè." The sheet bears also two papers with notes by Francis Boott and the penciled note of Dr. C. W. Swan: "The original specimen found by B. D. Greene at Tewksbury, Mass." Boott's comments are interesting though of no importance in settling the exact station: "These 3 plants I part with as a miser does gold entrusted to him. The Georgia one I suppose to be *C. Floridana*. . . . *C. fulva* & *C. laevigata* we owe to you & it behoves you to find them again. I take them to be *C. Greeniana* & *binervis* of Torrey [by which he obviously meant Dewey]."

The statement on Greene's label, that the plant grew with *Carex*

<sup>1</sup> Boott, Ill. Car. iv. 138 (1867).



*limosa* in a pond (presumably at Tewksbury) gives an important clue, for Tewksbury was one of B. D. Greene's great collecting grounds and he made its ponds and those of Andover famous. There he collected the types of *Juncus militaris* and *Utricularia resupinata*; and on the sandy and peaty margin of Round Pond are found many isolated northern colonies of Coastal Plain species: *Sagittaria Engelmanniana*, *Fuirena squarrosa*, *Xyris Smalliana*, etc. Round Pond has therefore been much visited and always furnishes good botanizing, even at the present time occasionally giving us new records. But the other pond of Tewksbury, Long Pond, is apparently little visited by botanists because its western side is bordered by an extensive quaking bog so deep and wet as to fatigue or drive away all but the most venturesome. Here, however, is the only known station in Middlesex County for *Carex limosa*! A specimen collected by B. D. Greene himself and labeled merely "Tewksbury" is in the Gray Herbarium, but material collected by William Boott on July 8, 1863 is marked "West side of Long Pond." No other station is cited in the *Middlesex Flora*, and it is obviously in this quaking bog or in the western margin of Long Pond, that we must seek the long-lost *C. Hornschuchiana* var. *laurentiana*. On June 17 last Mr. F. F. Forbes and the writer made a first attempt at rediscovery; but we were much too early, though we found young *C. limosa* in abundance. The traveling is difficult, consisting of waist-deep wallowing and tumbling, but by mid-July, when the prize *Carex* of the bog should be sought, traveling there may prove less formidable. At any rate the search is worth making and no effort should be spared to prove whether or not the almost forgotten plant still persists at Long Pond.

In view of the established occurrence of *Carex Hornschuchiana*, var. *laurentiana* in Newfoundland and its great rarity on our continent (found only at one of the ponds in Tewksbury) it is noteworthy that Haggett's Pond in Andover (lying just to the north of Long Pond) furnishes a case of somewhat similar distribution. By all means the most generally distributed grass upon Newfoundland is *Calamagrostis Pickeringii* and, though it is found locally upon our granitic mountains, it is known as a coastwise species only from Newfoundland, Cape Breton Island, and at a restricted station near the northern end of Haggett's Pond in Andover, a famous station discovered by Mr. John Robinson in 1879. It seems probable, then, that *Carex Hornschuchiana*, var. *laurentiana* like *Calamagrostis Pickeringii*, a plant

which at present finds its great development in the Newfoundland area but retains a tenuous hold at remote points in the line of its northward migration, is to be sought in Cape Breton and upon the mountains of northern New England and northern New York.

GRAY HERBARIUM.

REPORTS ON THE FLORA OF THE BOSTON  
DISTRICT,— XIII.

CYPERACEAE.

CAREX, D TO K.

**C. debilis** Michx., var. **Rudgei** Bailey. Woods and swamps, common.

**C. debilis** Michx., var. **Rudgei** Bailey  $\times$  **virescens** Muhl. Grove, Revere (*E. & C. E. Faxon*, June 10 to July 16, 1879–80; *H. A. Young*, June 17, 1879); Simpson Meadow, Campus, Wellesley (*K. M. Wiegand*, June 16, 1909).

**C. deflexa** Hornem. "Essex, Mass., . . . scarce," (Gray's Manual, 6th ed. 611, 1889, under var. *Deanei* Bailey); Blue Hill, Milton (*G. G. Kennedy*, May 27, 1910 and later).

**C. Deweyana** Schwein. Great Pastures, Salem (*J. H. Sears*, June 26, 1881); Roxbury (*J. A. Lowell*, June, 1846); Concord (*E. S. Hoar*, specimen in herb. of, according to Dame & Collins, Fl. Middlesex Co. 120, 1888).

**C. diandra** Schrank. Border of salt marsh, common, Newburyport (*F. F. Forbes*, June 14, 1903); Danvers (*J. H. Sears*, 1878); Beverly (*J. Robinson*, June 17, 1879); Boxford (*E. & C. E. Faxon*, June 20, 1878); bog, Lynnfield (*M. L. Fernald*, June, 1907); shore of Mud Pond, Natick (*K. M. Wiegand*, June 10, 1908); Wigwam Pond, Dedham (*C. E. Faxon*, May 26, 1878).

**C. diandra** Schrank, var. **ramosa** (Boott) Fernald. In brackish marsh, Newburyport (*F. F. Forbes*, June 13, 1903); Lexington (*E. Tuckerman*, no date); "wet meadow, head of pond, Fairy Land," Concord (*H. D. Thoreau*, June 30, —); open peat-bog, Sudbury



(*M. L. Fernald & H. M. Noyes*, June 14, 1901; *M. L. Fernald, W. P. Rich & E. F. Williams*, June 17, 1902).

**C. digitalis** Willd. Dry and moist woods. Not reported from northwestern and southeastern sections, but frequent elsewhere (24 stations reported).

[*CAREX EBURNEA* Boott. In the herbarium of the New England Botanical Club is a *Carex* labelled, "*Carex eburnea* Boott, Reading, Mass., legit W. H. Manning, July 14, 1882." Accompanying the specimen is another label with the above name in Wm. Boott's handwriting. This specimen is the authority for the record in Dame & Collins, Fl. Middlesex Co., 118, 1888, "Reading (W. H. Manning.) Rare." The specimen is not *Carex eburnea*, as it lacks the pale tubular sheaths characteristic of the species, and its identification is uncertain, especially as the perigynia are wanting. As there is a specimen of true *C. eburnea* from Vermont in the Manning Herbarium, now deposited in the herbarium of the New England Botanical Club, and as Mr. Boott would never have misnamed the Reading plant, it is plain that there has been a misplacement of specimen and label in this case.]

**C. exilis** Dewey. Bogs and wet meadows; nineteen stations in Essex and Middlesex counties.

Originally described in Silliman's Journal XIV, 351, 1828, Appendix to Caricography. After the original Latin description is the following note: "Flowers in May; found at Danvers, Mass., by Wm. Oakes, Esq., and is a singular and beautiful plant. It grows in small tufts, and also separate, in wet, cold meadows." On the same page Dewey describes a var. *squamacea* with the following note: "Flowers in May-June; found at Ipswich, Mass., by Mr. Oakes in dense and large tufts on the overflowed lands of Ipswich river."

**C. festucacea** Schkuhr. Sandy soil, filled land off Ipswich St., Boston, not abundant (*F. F. Forbes*, June 24, 1904); railroad by Charles River, Needham (*J. R. Churchill*, June 17, 1887); sandy plain near Lake Cochituate, Natick, not abundant, growing near *C. Bicknellii* with intermediates between the two (*F. F. Forbes*, June 24, 1906); dry railroad bank, Walpole (*G. G. Kennedy*, July 20, 1899); moist field, Mansfield (*J. A. Cushman*, June 1, 1908).

**C. festucacea** Schkuhr, var. **brevior** (Dewey) Fernald. Dry soil; many reports from central part of district, none from northern part, and only one (Norfolk) from southern part.

**C. filiformis** L. Bogs and wet places, common.

**C. flava** L. Wet places, occasional in northern section.

**C. flava** L., var. **elatior** Schlecht. Georgetown, Lynnfield, Malden, Cambridge, Wilmington, Lexington, Chelmsford, Concord and Dedham.

**C. flava** L., var. **rectirostra** Gaudin. No reports from northwest or southeast; well distributed elsewhere.

**C. foenea** Willd. Open woods; reports from scattered stations in central and northeastern portions.

**C. foenea** Willd., var. **perplexa** Bailey. Central and northeastern portions, occasional.

**C. folliculata** L. Swamps and wet woods; very common throughout.

**C. glaucodea** Tuckerm. Moist soil, south side of Blue Hill, Milton (*G. G. Kennedy*, June 14, 1901; other collections since to June 9, 1908).

**C. Goodenowii** J. Gay. Swamps and wet, sometimes brackish places; common throughout, especially near the coast.

**C. gracillima** Schwein. Deciduous woods and meadows. Not reported south of the Blue Hills, but common elsewhere.

**C. gracillima** Schwein., var. **humilis** Bailey. Revere (*H. A. Young*, June, 1879); Edgeworth, Malden (*Wm. Boott*, June 30, 1864).

**C. GRAYII** Carey. Waste grounds, Cambridge (*W. Deane*, June 27, 1887). Adventive from further west.

**C. grisea** Wahlenb. Readville, Hyde Park (*S. T. Olney*, May 29, 1868).

**C. HELODES** Link (*C. laevigata* Smith). Near Boston, perhaps at Tewksbury (*B. D. Greene*, prior to 1836). Not since collected and presumably a casual or adventive plant. See notes by M. L. Fernald, RHODORA, XIII, 243-248.

**C. HIRTA** L. Groves and fields; Danvers, Melrose, Medford, Newton, Ashland; Cambridge (*L. H. Bailey*, according to Dame & Collins, Fl. Middlesex Co., 115, 1888). No reports more recent than 1887.

**C. hormathodes** Fernald. Salt and fresh marshes and other wet places; frequent in eastern part of district, especially on the coast.

**C. hormathodes** Fernald, var. **invisa** (W. Boott) Fernald. Originally described by William Boott as *C. straminea* var. *invisa*, "Common on margins of ponds and in ditches in sandy soil, E.



Massachusetts. Mt. Desert, Maine." Boott's Massachusetts specimens were from Mystic Pond, Medford, "Ice Pond," Medford, and "Brookline branch R. R." The variety is apparently less common than typical *C. hormathodes*.

**C. hormathodes** Fernald, var. **Richii** Fernald. Frequent in central half of district, as far north as Ipswich, and south as Walpole. Named from specimens collected in Stoneham by W. P. Rich.

**C. Hornschuchiana** Hoppe, var. **laurentiana** Fernald & Wiegand. With *C. limosa*, western edge of Long Pond, Tewksbury (*B. D. Greene*, prior to 1836). Not since collected. See notes by M. L. Fernald, *RHODORA*, XIII, 243-248.

**C. hystericina** Muhl. Newburyport (*A. A. Eaton*, June 30, 1896); Beverly (*J. A. Lowell*, 1847); wet place, aqueduct near Benvenue St., Wellesley (*K. M. Wiegand*, July 6, 1908).

**C. hystericina** Muhl., var. **Cooleyi** Dewey. Fresh Pond, Cambridge (*Wm. Boott*, no date).

**C. intumescens** Rudge. Low ground, swamps and meadows; common throughout.

**C. intumescens** Rudge, var. **Fernaldii** Bailey. Wet woods, Arlington (*C. H. Knowlton*, June 22, 1907); wet place on bank of Charles River, Wellesley (*K. M. Wiegand*, Oct. 7, 1907); Heard's Pond, Wayland (*K. M. Wiegand*, July 17, 1908); wet field, south side of Blue Hill (*Wm. Boott*, July 7, 1853); rich woods on bank of Sudbury River (*A. J. Eames*, July 27, 1909).

C. H. KNOWLTON	} Committee on Local Flora.
J. A. CUSHMAN	
WALTER DEANE	
A. K. HARRISON	

## SALIX CALCICOLA, A LITTLE KNOWN NORTHERN WILLOW.

M. L. FERNALD AND K. M. WIEGAND.

**Salix calcicola**, n. sp. *S. Richardsoni*, var. *Macouniana* Bebb, Bot. Gaz. xiv. 50, t. 9 (1889). *S. lanata*, var. *Macouniana* Bebb, according to Macoun, Ann. Rep. Geol. Surv. Can., n. s. iii. 70 J (1889), *nomen nudum*.

Since this remarkable shrub was described by Mr. Bebb from one of the Twin Islands in James Bay,<sup>1</sup> it has been found on calcareous cliffs of Table-top Mountain, Gaspé County, Quebec, by *Fernald & Collins* (no. 211, August 7, 1906), on the shores of Kangalaksiorvik Bay, Labrador by *Owen Bryant* (no. 75, September, 1908) at Burwell, Hudson Strait and Churchill, Hudson Bay, lat. 58° 50' by *J. M. Macoun* (nos. 79,155 and 79,154, July 18 and August 3, 1910), and on limestone barrens at sea-level by Ingornachoix Bay and on Pointe Riche, Newfoundland, by *Fernald & Wiegand* (nos. 3151 and 3152, August 1 and 4, 1910). All this material is essentially uniform in character, agreeing with the original plate of *Salix Richardsoni*, var. *Macouniana* in its short-ovate to orbicular apiculate round-based or subcordate leaves, its dense aments and undivided stigmas. In these characters the eastern shrub is strongly contrasted with Hooker's *S. Richardsoni*, which has the narrowly obovate gradually acute leaves tapering at base, and the stigmas distinctly cleft. Besides these characters, already noted by Bebb, we find that in *S. calcicola* the midrib of the leaf is much wider and more conspicuous and the shorter petiole much broader than in *S. Richardsoni*. In the latter species, the stipules, as shown in Hooker's plate (Fl. Bor. Am. t. clxxxii) and in two authentic specimens in the Gray Herbarium, are lanceolate and bear three conspicuous gland-tipped teeth on each side. In *S. calcicola* the stipules are oblong-ovate to suborbicular, with numerous very fine glandular teeth. In all these characters the eastern shrub is so constant and distinctive that we feel no hesitation in raising it to specific rank; but since the name *Salix Macounii* has already been used in the genus it is advisable, in order to avoid possible confusion, to give the plant a new name.

*Salix calcicola* is quickly distinguished from all the other species known to us from eastern America by its large terminal (as well as rarely scattered) catkins and by the very large terminal hairy winterbuds which, when fully grown are 5-10 mm. in diameter. The fully mature leaves become quite glabrate and thick, varying from 2 to 5 cm. in length.

*Salix lanata* L. of Arctic Europe has been reported from "eastern

<sup>1</sup> In his account of the expedition to James Bay, Mr. James M. Macoun (Bot. Gaz. xiii. 117) says that "on the north Twin Island specimens of a new willow were collected which has been described by Mr. Bebb"; but, in the original publication of the plant, Bebb said "South Twin, James Bay. Collected July 17, 1887, by Mr. James M. Macoun, for whom it is named."



British America north of the Arctic circle and Greenland (Hook. Arc. Pl.)”<sup>1</sup> but Bebb remarks: “I have not seen specimens.”<sup>1</sup> It is not improbable that Hooker’s British American plant was *S. calcicola*, from which *S. lanata* is distinguished by its more elongate and pubescent leaves, its longer aments with more colored hairs, and by the 2-cleft stigmas.

---

STUMP-HEALING IN *PINUS STROBUS*.—A few years ago I noticed a white pine stump which was healing over! This was contrary to all experience. For such growth is uncommon or rare on low stumps even among our dicotyledons which ‘sucker up’ freely. And *Pinus Strobus* never grows again from the stump,—*P. rigida* being our only conifer that behaves in this manner. The tree had been cut about four years apparently, and during this time the wound-tissue had spread inward irregularly over the cut surface, in places nearly an inch. How could a leafless plant do this? But one answer was possible,—that nourishment was being derived from some other tree, and root-grafting was suspected as the means, though such could not be proved at the time. Later a similar growth was noted on a small hemlock stump in the Arnold Arboretum. This summer another case has come to my attention, and has supplied evidence showing that root-grafting is indeed the cause of these unusual growths. One pine among a group was cut, and the stump healed slowly for two or three years before dying. These trees stand on a gravelly knoll and their roots are nearly all exposed for some distance from the trunks. Grafting frequently occurred where crossings were formed, especially along a path where the roots had been wounded. It is probable that a parasitism of this sort is not uncommon among our trees, but the sight of an apparently lifeless pine-stump magically healing itself is almost startling.—ARTHUR J. EAMES, Cambridge, Massachusetts.

---

BOTANICAL LECTURES AT THE LOWELL INSTITUTE.—The Lowell Institute announces among its courses of free public lectures a series on Local Natural History to be given under the auspices of the Boston Society of Natural History. The series upon “The Wild Plants of Eastern Massachusetts” is to be given by Professor M. L. Fernald on

<sup>1</sup> Bebb, Bot. Gaz. xlv. 49 (1889).

Thursdays in January at 4 p. m., in the lecture-room of the Boston Society of Natural History. The dates and subjects are as follows:—

January 4. The Indigenous Flora of Eastern Massachusetts.

January 11. Some Effects of Civilization upon our Flora.

January 18. Some Local Problems in Geographic Botany.

January 25. The Probable Geographic Origin of our Flora.

Tickets may be secured, free of charge, at the Boston Society of Natural History; at the office of the Lowell Institute, 491 Boylston St.; or by sending a request, *accompanied by an addressed, stamped envelope*, to the Curator of the Lowell Institute, 491 Boylston St., Boston.

### ERRATA.

Page 76, line 6; for *oxycanthoides* read *oxyacanthoides*.

“ 89, “ 16; for *Thlapsi* read *Thlaspi*.

“ 89, “ 17; for *Loesellii* read *Loeselii*.

“ 89, “ 27; for *Siderites* read *Sideritis*.

“ 108, “ 28; for *K.* read *H.*

“ 109, “ 12; for *Altantic* read *Atlantic*.

“ 110, “ 7; for *our* read *one*.

“ 118, “ 19; for *Mr.* read *Mt.*

“ 123, “ 22; for *vegetation* read *vegetation*.

“ 124, “ 7; for *voilet* read *violet*.

“ 130, “ 3; for *carcina* read *caricina*.

“ 132, “ 25; *Insert comma after* for.

“ 138, “ 27; for *meliciodes* read *melicoides*.

“ 138, “ 38; *after p. insert* 131.

“ 143, “ 25; for *are* read *is*.

“ 145, “ 34; for *note 1* see *note 1* on page 146.

“ 145, “ 37; *insert* <sup>1</sup> *before* Warming.

“ 146, “ 30; for *Anticosti*,<sup>3</sup> read *Anticosti*,<sup>2</sup>

“ 146, “ 33; for *Lawrence*,<sup>1</sup> read *Lawrence*.<sup>3</sup>

“ 151, “ 30; for *note 1* see *note 1* on page 152.

“ 165, “ page heading; for *arundalana* read *arundelana*.

“ 186, “ 5; for *Acrochaete* read *Acrochaetium*.

“ 223, “ 18; for *tradactylites* read *tridactylites*.

*Vol. 13, no. 155, including pages 225 to 240 and one unnumbered portrait was issued 25 November, 1911.*



# INDEX TO VOLUME 13.

New scientific names are printed in full face type.

- Abies balsamea*, 138.  
*Acer circinatum*, 190; *rubrum*, 91, 139; *Saccharum*, 141, 148.  
*Acnida tuberculata*, 89.  
*Acrochaetium*, 186; *Dasyae*, 186; *Dictyotae*, 186; *Dufourii*, 187; *flexuosum*, 186; *Hoytii*, 186; *minimum*, 186.  
*Actaea rubra*, 125, 126.  
 Additions to the Description of *Streptopus longipes* Fernald, 237.  
*Adiantum*, 134, 141; *pedatum*, 141, 147, 161, var. *aleuticum*, 118, 132, 137, 141.  
*Agarics*, 57.  
*Agrimonia*, 18.  
*Agropyron caninum*, 89; *pungens*, 178; *tenerum*, 89.  
*Agrostis alba*, var. *aristata*, 177; *canina*, 30, 70.  
*Ahnfeldtia plicata*, 187, forma ***furcellata***, 187.  
*Aigeiros deltoides*, 12, 199; *virginiana*, 12, 198.  
*Alchemilla*, 117, 124; *officinalis*, 110.  
*Aletris farinosa*, 179.  
*Algae*, Notes on,—X, 184.  
*Alisma Plantago-aquatica*, 141.  
*Alliaria officinalis*, 89.  
*Alnus alnobetula*, 216; *maritima*, 220; *mollis*, 216; *rugosa*, 91; *viridis*, 216.  
*Amanita*, 57, 60, 62; *cariosa*, 60, 61, 66; *spissa*, 61.  
*Amanitopsis strangulata*, 61.  
*Amaranthus blitoides*, 89.  
*Amateur botanical Illustrating*, 93.  
*Amblystegium riparium*, 44; *vacillans* 44.  
*Ambrosia psilostachya*, 33.  
*Amelanchier oligocarpa*, 138.  
*America*, *Arrhenatherum elatius*,  **$\beta$**  *tuberosum* in, 207; *Cornus canadensis*, var. *intermedia* in eastern, 107; *Erucastrum Pollichii*, adventive in, 10; The Variations of *Lathyrus palustris* in eastern, 47.  
*Ammophila arenaria*, 128, 135, 140.  
*Anacamptodon splachnoides*, 45, 46.  
*Anaphalis margaritacea*, 25, 26, var. *occidentalis*, 25–27; Some boreal Species and Varieties of *Antennaria* and, 23.  
*Andromeda*, 114; *glaucophylla*, 54, 217, 220; *polifolia*, 217, 220.  
*Anemone*, 18; *canadensis*, 141; *multifida*, 148; *narcissiflora*, 222, 223; *parviflora*, 117, 129, 223; *riparia*, 141, 148, 161; *virginiana*, 218.  
*Angelica atropurpurea*, 123.  
 Another Station for *Centaurea diffusa*, 100.  
*Antennaria*, 23, 129; *alpina*, 24, 25, 127, var. ***cana***, 24, 25, 127, 129, 137; and *Anaphalis*, Some boreal Species and Varieties of, 23; *canadensis*, 23, 25; *carpatica*, 24, var. *humilis*, 24; ***eucosma***, 23, 24, 129, 130, 138; *lanata*, 24; *neodioica*, 23, 25, 139, var. *gaspensis*, 71; *Parlinii*, 25; *petaloidea*, 23; *plantaginifolia*, 24, 25; *pulcherrima*, 24.  
*Anthemis arvensis*, var. *agrestis*, 18.  
*Anthyllis*, 240; *Vulneraria*, 240; at Cromwell, Connecticut, 240.  
*Apocynum androsaemifolium*, 18, 139; *cannabinum*, 148.  
 Apple, Baked, 123.  
*Aquilegia*, 18.  
*Arabis*, 37; *albida*, 123; *alpina*, 119, 123; *brachycarpa*, 148; *Drummondii*, 37; *Holboellii*, 219; *laevigata*, 37, 38; A new, 37; *virginica*, 91; ***viridis***, 38.  
*Aralia*, 18; *hispida*, 139, 146, 149.  
*Arbor Vitae*, 115, 142.  
*Arctium*, 18.  
*Arctostaphylos alpina*, 117, 123, 136.  
*Arenaria*, 180; *arctica*, 118, 134, 136; *ciliata*, 118, var. *humifusa*, 118, 134; *groenlandica*, 180; *juni-perina*, 117; *litorea*, 117; *norvegica*, 118; *peplodes*, var. *robusta*, 135, 140; *verna*, var. *propinqua*, 117.  
*Arethusa*, 113, 114; *bulbosa*, 139.  
*Arisaema*, 18; *Dracontium*, 78; *triphyllum*, 78, 141.

- Armeria*, 216.  
*Arnica chionopappa*, 148; *gaspensis*, 219; *mollis*, 148.  
*Arrhenatherum bulbosum*, 208; *elatus*, 208, var. *tuberosum*, 207, in America, 207.  
*Artemisia Absinthium*, 90; *annua*, 70, 90; *biennis*, 90; *campestris*, var. *pubescens*, 90; *Dracunculus*, 90; *frigida*, 90; *pontica*, 71.  
*Asclepias*, 19; *amplexicaulis*, 19; *syriaca*, 18, 148.  
*Ascyrum hypericoides*, 151.  
 Ash, 57; Mountain, 238.  
*Aspidium aculeatum*, var. *Braunii*, 218; *Filix-mas*, 111; *Goldianum*, 238; *marginale*, 111; *simulatum*, 78, 177.  
*Asplenium angustifolium*, 238; *thelypteroides*, 111.  
*Aster*, 123, 142; *acuminatus*, 141, 148; *cordifolius*, 148; *juncus*, 221; *Lindleyanus*, 183; *longifolius*, 33; *Lowricanus*, 32, var. *lanceolatus*, 32; *macrophyllus*, 141, 142, 148, 161; *nemoralis*, 135, 139; *radula*, 135, var. *strictus*, 96; *spectabilis*, 8; *subulatus*, 150; *tardiflorus*, 183; *umbellatus*, 139.  
*Astragalus alpinus*, 218; *frigidus*, var. *americanus*, 148.  
*Atriplex patula*, var. *hastata*, 89, 125, 180; var. *littoralis*, 180.  
*Avena bulbosa*, 208; *elatior*, var. *tuberosa*, 208; *tuberosa*, 207.  
*Baccharis halimifolia*, 243.  
 Baked Apple, 123.  
 Balm of Gilead, 237.  
*Barbarea orthoceras*, 123.  
*Barbula fallax*, 46.  
 Barnes, C. R., [Notice of Work], 16.  
 Barnstable, Massachusetts, *Cynanchum nigrum* in, 70.  
 Bartlett, H. H., *Euphorbia arundeliana*, an Ally of *Euphorbia Ipecacuanhae*, 163; On Gynodioecism in *Plantago lanceolata*, 199; *Populus virginiana* and *P. angulata*, 12; *Ptelea mollis*, var. *cryptoneura*, a Wafer-ash of the Georgia Sand Hills, 80; Systematic Studies on *Oenothera*,—I. *Oenothera Tracyi*, sp. nov., 209.  
 Bartlett's *Dioscoreae* of the United States, 34.  
*Bartonia iodandra*, 116, 159; *virginica*, 182.  
 Bearberry, 123.  
 Bedstraw, 237.  
 Beeches, 86.  
 Berberis, 18.  
 Berry, Partridge, 216.  
*Berteroa incana*, 89.  
*Betula*, 113, 114; *alba*, var. *cordifolia*, 238; *glandulosa*, 123, 136, 219; *lenta*, 111, 206, 207; *lutea*, 46, 206, 207; *nana*, var. *Michauxii*, 113, 116; *nigra*, 151, 155; *pumila*, 138, 217.  
*Bidens cernua*, 141; *hyperborea*, 183.  
 Birch, 57, 113; Black, 206, 207; Cherry, 206; Gray, 207; Paper, 207; Silver, 207; Sweet, 206; to be restricted, The Range of the Black, 206; Yellow, 206, 207.  
 Bissell, C. H., A Day at Congamond Lakes, 53; Notes on Connecticut Plants, 30.  
 Black Birch, 206, 207, to be restricted, Range of the, 206; Poplar, 195; Poplar, Virginian, 198; Spruce, 114.  
 Blake, S. F., Pteridophyte Notes from eastern Massachusetts, 101; A *Scirpus* new to New Hampshire, 55.  
 Blanchard, W. H., *Lycopodium complanatum* near Hartland, Vermont, 211; *Lycopodium flabelliforme*, 168; A necessary Change of Name, 55; A new Variety of *Rubus canadensis*, 193; The Range of the Black Birch to be restricted, 206.  
 Blewitt, A. E., Some introduced Plants of Connecticut, 88.  
 Blueberries, 86.  
*Boletinus cavipes*, 57; *paluster*, 57.  
*Boletus*, 57; *clintonianus*, 57; *eximius*, 57; *punctipes*, 57; *spectabilis*, 57.  
 Boreal Variety of *Fragaria virginiana*, 106.  
 Boston District, Reports on the Flora of the,—VIII, 27, IX, 72, X, 82, XI, 104, XII, 232, XIII, 248.  
 Botanical Club, Notes from the Phaenogamic Herbarium of the New England,—II, 177; Club, Sixteenth annual Meeting of the Vermont, 237; Club, The winter Meeting of the Vermont, 87; Expedition to Newfoundland and southern Labrador, 109; Lectures at the Lowell Institute, 253; Society of Maine, Winter Meeting of the Josselyn, 90.  
 Botanizing in central Connecticut, 77.

- Botany, A new Textbook on, 16.
- Botrychium*, 14, 15; from Tennessee, Notes on, 14; *Lunaria*, 119, 147; *obliquum*, 14, 103, var. *dissectum*, 15, var. *oneidense*, 102; *ternatum*, var. *intermedium*, 103; *virginianum*, 15, 125.
- Brainerd, E., Cyrus Guernsey Pringle 225.
- Brassica Erucastrum*, var. *ochroleuca*, 11; *juncea*, 10, 89; *obtusangula*, var. *Pollichii*, 11; *ochroleuca*, 11.
- Bromus*, 88; *inermis*, 88; *squarrosus*, 88; *tectorum*, 88.
- Bush, B. F., The Missouri Rhexias, 166.
- Calamagrostis hyperborea*, 136; *ladorica*, 119; *lapponica*, 123; *neglecta*, 123; *Pickeringii*, 116, 131, 247; *purpurascens*, 219.
- Calla palustris*, 78, 220.
- Calliargon trifarium*, 44, 45.
- Callithamnion Lenormandi*, 186, 187.
- Callitriche anceps*, 133.
- Calopogon pulchellus*, 128, 139, 220.
- Calothrix*, 185; *Braunii*, 185; *crustacea*, 185; *Kawraiskyi*, 185.
- Caltha palustris*, 123.
- Calypso*, 221; *borealis*, 221; *bulbosa*, 115, 161, 221.
- Camelina microcarpa*, 89.
- Campanula rotundifolia*, var. *alaskana*, 127.
- Camptothecium nitens*, 46.
- Cantharellus dichotomus*, 63.
- Cape Cod; third Note, Flora of Lower, 17.
- Capsella Bursa-pastoris*, 91; *elliptica*, 137.
- Carduus*, 238; *leiocephalus*, 240; *plattensis*, 240; *spinosissimus*, 239.
- Carex*, 130, 233, 247, 248; *acutiformis*, 233; *acenea*, 178; *aestivalis*, 30, 233; *alata*, 233; *albolutescens*, 233, var. *cumulata*, 233; *aquatilis*, 233; *arctata*, 233; *atlantica*, 135; *aurea*, 233; *Bicknellii*, 233; *bicolor*, 127; *binervis*, 243, 244; *bromoides*, 233; *brunnea*, 233; *brunnescens*, 233; *bullata*, 234, var. *Greenii*, 234, var. *Greenii* × *lupulina*, 234, var. *Greenii* × *vesicaria*, 234, var. *Greenii* × *vesicaria*, var. *monile*, 234; *canescens*, var. *disjuncta*, 234, var. *sublobliacea*, 234, var. *vulgaris*, 234; *capillaris*, var. *elongata*, 125; *caryophyllea*, 234; *castanea*, 115, 138; *cephaloidea*, 69, 234; *cephalophora*, 234; *communis*, 234; *comosa*, 234; *conoidea*, 234; *Crawei*, 30; *Crawfordii*, 234; *crinita*, 234, var. *gynandra*, 235, var. *minor*, 235; *cristata*, 235; *debilis*, var. *Rudgei*, 69, 117, 248, var. *Rudgei* × *virescens*, 69, 248; *deflexa*, 117, 248, var. *Deanei*, 248; *Deweyana*, 125, 126, 248; *diandra*, 248, var. *ramosa*, 248; *digitalis*, 249; *eburnea*, 117, 138, 249; *exilis*, 249, var. *squamacea*, 249; *festucacea*, 249, var. *brevior*, 249; *filiformis*, 250; *flava*, 130, 250, var. *elator*, 250, var. *gaspensis*, 132, 138, var. *rectirostra*, 250; *floridana*, 246; *foenea*, 250; var. *perplexa*, 250; *folliculata*, 148, 250; *fulva*, 130, 131, 243, 244, 245, 246, *fulvescens*, 245; *glaucodea*, 250; *Goodenowii*, 250; *gracillima*, 250, var. *humilis*, 250; *Grayii*, 250; *Greeniana*, 130, 243-245; *grisea*, 250; *gynocrates*, 115; *helodes*, 243-246, 250; *hirta*, 250; *hormathodes*, 135, 140, 250, var. *invisa*, 128, 250, var. *Richii*, 251; *Horns chuchiana*, 130, 131, 243-245, var. *laurentiana*, 130, 138, 245-247, 251; *hystericina*, 251, var. *Cooleyi*, 251; *intumescens*, 117, 149, 251, var. *Fernaldii*, 251; *laevigata*, 243, 244, 245, 250; *lagopina*, 219; *laxiflora*, var. *leptonervia*, 125; *limosa*, 116, 247, 251; *lurida*, 141, 148; *Michauxiana*, 138; *novae-angliae*, 30; *oligosperma*, 217, 218; *panicea*, 243; *pauciflora*, 95; *pedata*, 129, 137; *pelva*, 244; *polygama*, 116; *prasina*, 178; *ptychocarpa*, 151; *rariflora*, 122, 136; *retrorsa*, 141, 148, 161; *rupestris*, 129; *scirpoidea*, 117, 238; *silicea*, 128, 135, 140; *sterilis*, 135, 140; *straminea*, var. *invisa*, 250; *stricta*, 141; *stylosa*, 122; *subulata*, 78, 79, 151; *trisperma*, 54, var. *Billingsii*, 116; 135; *umbellata*, 69, var. *brevirostris*, 68, 69; *vaginata*, 115, 125; *vestita*, 179; *virescens*, 69; *vulpinoidea*, 139, 146.
- Carices of eastern Massachusetts, Two lost, 243.
- Carya*, 91; 217.
- Cassia Chamaecrista*, 8; *mari-landica*, 34.



- Cassiope tetragona, 223.  
 Castalia odorata, 149, 191.  
 Castanea, 90, 217; dentata, 219, 220; pumila, 90, in eastern Massachusetts, 90.  
 Catabrosa aquatica, 122.  
 Cedar, 57.  
 Centaurea diffusa, 100, Another Station for, 100.  
 Cerastium, 216; alpinum, 124.  
 Ceratophyllum, 79.  
 Chaetochloa, 174-176; occidentalis, 175; versicolor, 175.  
 Chamaedaphne, 215.  
 Chamaeraphis, 174.  
 Chantransia, 186; **Dasyae**, 186; **Dictyotae**, 186; Dufourii, 187; **flexuosa**, 186; **Hoytii**, 186; **minima**, 186.  
 Chase, A., Amateur botanical Illustrating, 93; Arrhenatherum elatius,  $\beta$  tuberosum in America, 207; Ornithological Observations on Cleistogamy, 76; The subterranean Organs of Cinna arundinacea, 9.  
 Chasea, 173-175; prolifera, 175; pubescens, 175; violacea, 175.  
 Chelone glabra, 146.  
 Chenopodium, 22; Botrys, 22; carinatum, 22; glaucum, 31; humile, 180; urbicum, 89.  
 Cherry Birch, 206.  
 Chestnut, 57.  
 Chimaphila maculata, 21; umbellata, 21, 147.  
 Chiogenes, 141; hispidula, 125.  
 Chrysophyllum protensum, 45.  
 Chrysopsis Nuttallii, 239; pilosa, 239.  
 Chrysosplenium americanum, 141.  
 Cicuta bulbifera, 128, 139, 149.  
 Cinna arundinacea, 9, 207, 208, The subterranean Organs of, 9; latifolia, 125.  
 Cinquefoil, 238.  
 Circaea alpina, 138.  
 Cirsium, 238, 239; **americanum**, 239; Drummondii, 240; **Eatonii**, 240; eriocephalum, var. leioccephalum, 240; foliosum, 240; horridulum, 239; muticum, 111, var. monticola, 111; Notes on the Genus, 238; palustre, 116; **plattense**, 240; pumilum, 111, 183; spinosissimum, 238, 239.  
 Cladium, 105; mariscoides, 105; reticulata, 91.  
 Clavaria ligula, 64; muscoides, 60; pallescens, 64, 65.  
 Claytonia caroliniana, 111.  
 Cleistogamy, Ornithological Observations on, 76.  
 Clematis verticillaris, 31, 148; virginiana, 141, 148.  
 Clemson College, South Carolina, Plants in Flower February 8, 1911, at, 91.  
 Clintonia, 220; borealis, 78, 125, 220.  
 Club, Notes from the Phaenogamic Herbarium of the New England Botanical,—II, 177; Sixteenth annual Meeting of the Vermont Botanical, 237; Winter Meeting of the Vermont Botanical, 87.  
 Cnicus, 238; americanus, 239; carlinoides, var. americanus, 239; spinosissimus, 239.  
 Cochlearia anglica, 119, 137, 223; officinalis, 124, 216; tridactylites, 223.  
 Collins, F. S., Flora of Lower Cape Cod; third Note, 17; Notes on Algae,—X, 184.  
 Collybia strictipes, 60, 65.  
 Comandra livida, 95.  
 Comment on the Use of the Term Labrador in Natural History, 236.  
 Compositae, 93.  
 Congamond Lakes, A Day at, 53.  
 Conioselinum chinense, 125.  
 Connecticut, Anthyllis Vulneraria at Cromwell, 240; Botanizing in central, 77; Mosses, Notes on,—II, 40; Observations on some Plants of eastern, 68; Plants, Notes on, 30; Some introduced Plants of, 88.  
 Convolvulus arvensis, 89; Sepium, 149.  
 Corallorrhiza maculata, 105; striata, 115.  
 Corallorrhiza maculata, 238; trifida, 78.  
 Corema, 181; Conradii, 135, 140, 149, 161, 181.  
 Coreopsis rosea, 8, 152.  
 Cornus alternifolia, 149; canadensis, 107, 108, 138, var. intermedia, 107, var. intermedia in eastern America, 107; circinata, 146; stolonifera, 216; suecica, 107, 108.  
 Correction regarding Professor Penhallow, 56.  
 Corydalis sempervirens, 141.  
 Corylus rostrata, 149.  
 Cotton-grass, 113, 123.  
 Coulter, J. M., [Notice of Work], 16.  
 Cowles, H. C., [Notice of Work], 16.

- Cranberry, 217; Mountain, 136.  
 ·Crataegus, 18; coccinea, 36.  
 Cromwell, Connecticut, Anthyllis  
   Vulneraria at, 240.  
 ·Crotalaria sagittalis, 8.  
 Cruciferae, 89.  
 ·Cryptogramma Stelleri, 137.  
 ·Curlew-berry, 123.  
 ·Currant, Skunk, 138.  
 ·Cuscuta arvensis, 8, 151; compacta,  
   8.  
 ·Cushman, J. A., Reports on the  
   Flora of the Boston District,—  
   VIII, 27, IX, 72, X, 82, XI, 104,  
   XII, 232, XIII, 248; Three addi-  
   tional Plants from Nantucket,  
   105.  
 ·Cynanchum nigrum, 70, 91, in  
   Barnstable, Massachusetts, 70.  
 ·Cynoglossum officinale, 89.  
 Cyperaceae, 5, 6, 27, 72, 83, 104, 233,  
   248.  
 ·Cyperus, 22, 27; aristatus, 27, 53;  
   dentatus, 28, 53, var. ctenostachys,  
   28; diandrus, 28; Engelmanni,  
   28; erythrorhizos, 28; esculentus,  
   28, 218; ferax, 28; filiculmis, 28,  
   var. macilentus, 28; fuscus, 28;  
   Grayii, 22, 27; Nuttallii, 29, 178;  
   ovularis, 29; rivularis, 29; strigo-  
   sus, 29, var. capitatus, 29, var.  
   compositus, 29, var. robustior, 29.  
 Cyripedium acaule, 97, 139; hir-  
   sutum, 114, 115, 132.  
 Cystopteris bulbifera, 124, 132; mon-  
   tana, 124, 219.  
 Dalibarda repens, 141, 161, 229.  
 Danforth, C. H., A Dimorphism in  
   Tiarella cordifolia, 192.  
 Danthonia, 134; intermedia, 118,  
   137.  
 Davis, S., Some fleshy Fungi of  
   Stow, Massachusetts, 57.  
 Day at Congamond Lakes, 53.  
 Deane, W., A Correction regarding  
   Professor Penhallow, 56; David  
   Pearce Penhallow, 1; Recent  
   Treatment of Panicum compared  
   with that of Gray's Manual, 66;  
   Reports on the Flora of the Boston  
   District,—VIII, 27, IX, 72, X,  
   82, XI, 104, XII, 232, XIII, 248;  
   Teratology in Trillium ovatum  
   Pursh, 189.  
 Dentaria, 229.  
 Dermocarpa, 185; **Vickersiae**, 184.  
 Deschampsia caespitosa, 131.  
 Desmodium, 18, 19, 21, 33; canes-  
   cens, 33; obtusum, 19.  
 Dewberry, 123.  
 Diapensia, 123; lapponica, 117, 136.  
 Dicksonia, 102; punctilobula, 101.  
 Dicranum, 41, 42; condensatum, 40,  
   41; Drummondii, 42; fulvum,  
   42; Muhlenbeckii, 41; pallidum,  
   40, 41; sabuletorum, 40, 41;  
   scoparium, 41; spurium, 40, 41,  
   var. condensatum, 40; strictum,  
   43; undulatum, 42; viride, 42.  
 Dictyopteris delicatula, 184.  
 Diervilla Lonicera, 117, 146.  
 Digitalis purpurea, 132.  
 Dimorphism in Tiarella cordifolia,  
   192.  
 Dioscorea, 34, 35; floridana, 35;  
   glauca, 35; hirticaulis, 35; pani-  
   culata, 35, var. glabrifolia, 35;  
   quaternata, 35; sativa, 35, 36;  
   villosa, 35, 36.  
 Dioscoreae of the United States,  
   Bartlett's, 34.  
 Diplotaxis, 11; bracteata, 11; mur-  
   alis, 11; tenuifolia, 11.  
 Dogwood, Poisonous, 57.  
 Draba, 216; arabisans, 117; aurea,  
   219; caroliniana, 8; incana, 119,  
   var. confusa, 119.  
 Dracocephalum parviflorum, 219;  
   thymiflorum, 212, a casual Plant  
   at Westford, Massachusetts, 212.  
 Drepanocladus scorpioides, 45.  
 Drosera anglica, 127, 133; linearis,  
   132; rotundifolia, 138.  
 Drummondia clavellata, 46.  
 Dryas Drummondii, 141, 148, 223;  
   integrifolia, 127, 129, 136, 219,  
   223, var. canescens, 129; octo-  
   petala, 223.  
 Dryopteris cristata, 116, 139, 146;  
   Filix-mas, 116, 137; fragrans, 111,  
   218, 219; marginalis, 141, 147.  
 Dulichium, 29; arundinaceum, 29.  
 Eames, A. J., Stump-healing in  
   Pinus Strobus, 253.  
 Echenias carlinoides, var. nutans,  
   239.  
 Echinochloa Walteri, 21.  
 Echinodorus tenellus, 151, 155.  
 Eelgrass, 215.  
 Elatine americana, 135, 140.  
 Eleocharis, 72; acicularis, 72; di-  
   andra, 178; Engelmanni, 72, var.  
   defonsa, 72; interstincta, 72, 151;  
   nitida, 134, 138; obtusa, 72, 141;

- olivacea, 72, 178; ovata, 72; palustris, 72, var. glaucescens, 72; pauciflorus, 221; quadrangulata, 72, 151; Robbinsii, 72, 73; rostrelata, 73, 178; tenuis, 73; Torreyana, 151; tricostata, 151; tuberculosa, 73.
- Elm, 57.
- Elodea, 54, 79.
- Elymus arenarius, 117, 122; brachystachys, 89; virginicus, 139.
- Empetrum nigrum, 95, 99, 123, 136, 216, var. purpureum, 117, 123.
- Entoloma, 58; clypeatum, 58, 59; grande, 59, 65; modestum, 59, 65; nidorosum, 63; rhodopolium, 63; sinuatum, 59; strictius, 58, 65.
- Epigaea, 193; repens, 97, 139, 193, 216.
- Epilobium, 123; angustifolium, 138; densum, 220; latifolium, 111, 132; lineare, 220; molle, 148; palustre, 188, var. **longirameum**, 188.
- Equisetum pratense, 115; scirpoides, 125; variegatum, 130.
- Eragrostis megastachya, 18, 88, 178; pilosa, 177.
- Ericaceae, 221.
- Erigeron, 124; acris, var. asteroides, 148, var. oligocephalus, 124; hysopifolius, 117; philadelphicus, 141, 148.
- Eriocaulon, 93; septangulare, 117, 135.
- Eriolytrum, 174.
- Eriophorum, 104, 113, 114; angustifolium, 123; callitrix, 104, 123, var. erubescens, 113, 116, 123; Chamissonis, 123, 147; gracile, 104, 123, 138; tenellum, 104, 123; vaginatum, 217; virginicum, 104, var. album, 104; viridi-carinatum, 104.
- Errata, 254.
- Erucastrium, 11; bracteatum, 11; inodorum, 11; Pollichii, 11, adventive in America, 10.
- Erythrotrichia ceramicola, 187.
- Eupatorium aromaticum, 8; leucolepis, 152; perfoliatum, 148; purpureum, 91, var. foliosum, 132, var. maculatum, 139; rotundifolium, 8; urticaefolium, 141, 148.
- Euphorbia, 93; **arundelana**, 163-165, an Ally of Euphorbia Ipecacuanhae, 163; corollata, 164; Helioscopia, 181; ipecacuanha, 163; Ipecacuanhae, 163, 164, Euphorbia arundelana an Ally of, 163; maculata, 18; Peplus, 181; polygonifolia, 181.
- Euphrasia, 110, 116, 128; Randii, 110.
- Euthamia floribunda, 106, 107.
- Eye Bright, 128.
- Fairy Flax, 116.
- Federation of Natural History Societies, New England, 187.
- Fern, Goldie's, 238.
- Fernald, M. L., A boreal Variety of *Fragaria virginiana*, 106; A botanical Expedition to Newfoundland and southern Labrador, 109; *Cornus canadensis*, var. *intermedia* in eastern America, 107; *Epilobium palustre*, var. *longirameum*, 188; Harshberger's Phytogeographic Survey of North America, 213; A new Species of *Scirpus* from Massachusetts and New Jersey, 4; The northern Variety of *Gaylussacia dumosa*, 95; Notes from the Phaenogamic Herbarium of the New England Botanical Club,—II, 177; A pubescent Variety of *Vaccinium vacillans*, 235; *Salix calcicola*, a little known northern Willow, 251; Some boreal Species and Varieties of *Antennaria* and *Anaphalis*, 23; Two lost Carices of eastern Massachusetts, 243; The Variations of *Lathyrus palustris* in eastern America, 47; The Varieties of *Ribes hirtellum*, 73.
- Festuca altaica, 118; ovina, var. brevifolia, 117, var. supina, subvar. pubiflora, 123; rubra, var. subvillosa, 70; scabrella, 118, 137.
- Fimbristylis, 73; Frankii, 73.
- Fink, B., [Notice of Work], 15.
- Flammula betulina, 64.
- Flax, Fairy, 116.
- Fletcher, E. F., *Dracocephalum thymiflorum* a casual Plant at Westford, Massachusetts, 212.
- Flora of Lower Cape Cod; third Note, 17; of the Boston District, Reports on the,—VIII, 27, IX, 72, X, 82, XI, 104, XII, 232, XIII, 248.
- Flynn, N. F., Sixteenth annual Meeting of the Vermont Botanical Club, 237; The winter Meeting of the Vermont Botanical Club, 87.
- Fragaria canadensis*, 106; *terracenovae*, 106; *vesca*, var. *americana*, 106; *virginiana*, 106, A boreal



- Variety of, 106, var. **terraenovae**, 106.  
*Fraxinus pennsylvanica*, 218.  
*Fuirena*, 104; *squarrosa*, 104, 247.  
*Funaria flavicans*, 43; *hygrometrica*, 43.  
 Fungi of Stow, Massachusetts, Some fleshy, 57.  
*Galium kamtschaticum*, 116, 138; *labradoricum*, 115, 119, 138; *triflorum*, 125; *verum*, 89, 237.  
 Gates, F. C., An Addition to the Description of *Streptopus longipes* Fernald, 237.  
*Gaultheria procumbens*, 97, 139.  
*Gaylussacia*, 96; *baccata*, 128, 139, 149, forma *leucocarpa*, 182; *dumosa*, 54, 96-99, The northern Variety of, 95, var. **Bigeloviana**, 99, 128, 135, 140, 149; *hirtella*, 97.  
*Gentiana linearis*, 182; *nesophila*, 119, 130, 138.  
 Georgia Sand-hills, *Ptelea mollis*, var. *cryptoneura*, a Wafer-ash of the, 80.  
*Geranium*, 18.  
*Gerardia parvifolia*, 151; *paupercula*, 19; *pedicularia*, 19; *purpurea*, 19; *virginica*, 182.  
*Geum*, 221; *macrophyllum*, 115, 125, 138.  
*Glaux maritima*, 105.  
*Glyceria acutiflora*, 92, in Vermont, 92; *borealis*, 92; *canadensis*, 139.  
*Gnaphalium decurrens*, 55; *polycephalum*, 148; *purpureum*, 33, 152.  
 Goldenrods, 123.  
 Goldie's Fern, 238.  
*Gomphidius*, 57; *gracilis*, 63; *maculatus*, 63; *roseus*, 60; *vinicolor*, 63, 65.  
*Goniotrichum elegans*, 187.  
*Goodyera pubescens*, 216, 218, 219; *tesselata*, 216.  
 Gramineae, 66.  
 Grass, Cotton, 123; Scurvy, 119, 124.  
 Gray Birch, 207.  
*Grindelia squarrosa*, 89.  
*Grossularia hirtella*, 73, 75; *inermis*, 74; *klamathensis*, 74.  
*Gynodioecism* in *Plantago lanceolata*, 199.  
*Habenaria blephariglottis*, 77, 128, 135; *dilatata*, 115, 138, 179; *fimbriata*, 77; *hyperborea*, 115, 138; *lacera*, 220; *macrophylla*, 179; *obtusata*, 138.  
*Halenia deflexa*, 138.  
 Hall, J. G., Plants in Flower February 8, 1911, at Clemson College, South Carolina, 91.  
 Harger, E. B., A new *Arabis*, 37.  
 Harrison, A. K., Reports on the Flora of the Boston District,—VIII, 27, IX, 72, X, 82, XI, 104, XII, 232, XIII, 248.  
 Harshberger's Phytogeographic Survey of North America, 213.  
 Hartland, Vermont, *Lycopodium complanatum* near, 211.  
*Hedeoma hispida*, 31.  
*Hedysarum alpinum*, 119, 129, var. *americanum*, 119; *americanum*, 119; *boreale*, 119, 218.  
*Helianthemum canadense*, 181; *major*, 181.  
*Helianthus divaricatus*, 18; *Maximiliani*, 89; *petiolaris*, 89; *subrhomboideus*, 103, in New Hampshire, 103.  
*Hemicarpha*, 53, 104; *micrantha*, 30, 104.  
 Hemlock, 57.  
*Hepatica*, 91; *triloba*, 91.  
*Heracleum lanatum*, 123, 138.  
 Herbarium, Notes from my, 71.  
*Hibiscus*, 21; *Moscheutos*, 21, 163.  
 Hickory, 57.  
*Hieracium*, 20; *canadense*, 20; *florentinum*, 219; *floribundum*, 219; *Greenii*, 20; *Gronovii*, 20; *marianum*, 20; *pratense*, 219; *praealtum*, 218, 219; *scabrum*, 20; *venosum*, 20.  
 Hill, E. J., Bartlett's *Dioscoreae* of the United States, 34.  
 Hitchcock, A. S., Type of the Genus *Panicum*, 173.  
*Holcus bulbosus*, 208.  
*Hordeum boreale*, 125, 127, 137; *jubatum*, 89.  
*Hottonia inflata*, 108, a winter Annual, 108.  
 Hubbard, F. C., *Anthyllis Vulneraria* at Cromwell, Connecticut, 240.  
 Huckleberry, 86.  
*Hudsonia ericoides*, 135, 140, 161, 181; *tomentosa*, 181.  
*Hydrocotyle umbellata*, 8.  
*Hygrophorus*, 57, 59, 66; *coloratus*, 64, 65; *flavodiscus*, 63, 65; *fuliginosus*, 63, 65; *hypothecus*, 63, 64; *laetus*, 59; *laurae*, 63, 65; *luridus*, 59.

- Hymenomycetes Europaei, 63.  
*Hypericum boreale*, 146; *virginicum*, 128, 146.  
*Hypnum*, 221.
- Ilex glabra*, 8, 150, 159, 161; *laevigata* 54; *opaca*, 151.  
 Important Publication on Lichens, 15.  
*Inocybe*, 57, 62; *commixta*, 62, 65; *echinata*, 60, 62; *geophylla*, 62; *intricata*, 62, 65.  
*Iris*, 110, 227; *setosa*, var. *canadensis*, 110; *verticillata*, 117, 139, 146, 149.  
*Iva xanthifolia*, 89.  
 Ivy, Poisonous, 57.  
*Ixophorus*, 174, 175.
- Josselyn Botanical Society of Maine, Winter Meeting of the, 90.  
*Juglans*, 217.  
*Juncus acuminatus*, 179; *alpinus*, var. *insignis*, 138; *aristulatus*, 151; *brachycarpus*, 151, 152; *canadensis*, 139; *compressus*, 138; *effusus*, 70, var. *compactus*, 70, var. *conglomeratus*, 31, 70, 140, var. *decipiens*, 179, var. *Pylaei*, 139; *militaris*, 247; *pelocarpus*, 128, 135; *stygius*, 124; *subtilis*, 128; *tenuis*, 146; *trifidus*, 117, 136, 238; *triglumis*, 124, 127, 130, 137.  
*Juniperus communis*, 215, var. *montana*, 132, 215; *horizontalis*, 117, 216; *Sabina*, 216, var. *procumbens*, 216; *virginiana*, 91, 216, 219, 220, var. *humilis*, 216.
- Kalmia angustifolia*, 54; *glauca*, 217; *latifolia*, 182.  
 Kidney Vetch, 240.  
 King Devil Weed, 219.  
 Kirk, G. L., *Glyceria acutiflora* in Vermont, 92.  
 Knowlton, C. H., *Cynanchum nigrum* in Barnstable, Massachusetts, 70; *Hottonia inflata* a winter Annual, 108; Notes on certain Leguminosae, 33; Reports on the Flora of the Boston District, — VIII, 27, IX, 72, X, 82, XI, 104, XII, 232, XIII, 248; Winter Meeting of the Josselyn Botanical Society of Maine, 90.  
*Kobresia caricina*, 124, 127, 130, 137; *elachycarpa*, 218.
- Labiatae, 93.
- Labrador, A botanical Expedition to Newfoundland and southern, 109; in Natural History, A Comment on the Use of the Term, 236.  
*Lactarius deliciosus*, 131.  
*Lactuca Scariola*, 10, 89, var. *integrata*, 89.  
 Ladies' Mantle, 124, Slipper, 114.  
*Lamium amplexicaule*, 91.  
 Larch, 114.  
*Larix laricina*, 137.  
*Lathyrus altaicus*, 52; *decaphyllus*, var. *minor*, 52; *myrtifolius*, 47, 48, 50, 52, var. *macranthus*, 48–50; *palustris*, 47–51, var. *genuinus*, 47, in eastern America, The Variations of, 47, var. *latifolius*, 47, var. *linearifolius*, 47–49, 51, var. **macranthus**, 50, var. *myrtifolius*, 47, 48, 52, var. *pilosus*, 48–51; *pilosus*, 48, 49, 51; *stipulaceus*, 52; *viciaeformis*, 47, 51.  
*Lechea intermedia*, 31; *maritima*, var. *interior*, 31.  
*Ledum latifolium*, 217.  
*Leersia oryzoides*, 148.  
 Leguminosae, 119; Notes on certain 33.  
 Leontodon, 18.  
*Lepidium apetalum*, 89.  
*Lepiota*, 62; *clypeolaria*, 60; *haematosperma*, 62; *metulisporea*, 60; *procera*, 61.  
*Leptochloa fascicularis*, 151.  
*Leskea*, 43; *gracilescens*, 43; *obscura*, 43, 44; *polycarpa*, 43, 44.  
*Lespedeza*, 33; *capitata*, 19, var. *velutina*, 33; *hirta*, var. *oblongifolia*, 31; *procumbens*, 19; *Stuvei*, 19.  
*Lesquerella*, 129; *arctica*, 223, var. *Purshii*, 127, 129–131, 138, 223.  
*Leucojum aestivum*, 179.  
 Lichens, 91; An important Publication on, 15.  
*Ligusticum scoticum*, 215, 216.  
*Lilium*, 227; *canadense*, 141, 161; *superbum*, 8.  
 Lily, Water, 191.  
*Limodorum tuberosum*, 220.  
*Limosella aquatica*, var. *tenuifolia*, 182.  
 Linnaea, 217; *borealis*, var. *americana*, 78, 125.  
*Linum catharticum*, 116; *floridanum*, 151.  
*Liriodendron*, 217; *tulipifera*, 217.  
*Listera auriculata*, 116, 138; *convallarioides*, 138; *cordata*, 138.

- Lithospermum arvense*, 89.  
 Liverwort, 221.  
*Lobelia Dortmanna*, 139; *Kalmii*, 138; *siphilitica*, 218.  
*Loiseleuria*, 123; *procumbens*, 117.  
*Lonicera canadensis*, 141, 161; *dioica*, 182; *involucrata*, 115; *oblongifolia*, 111, 217, 221.  
*Lophotocarpus spongiosus*, 177.  
 Lowell Institute, Botanical Lectures at the, 253.  
*Ludvigia sphaerocarpa*, 8.  
*Luzula spicata*, 123.  
*Lychnis*, 134; *alpina*, 118, 136.  
*Lycopodium alopecuroides*, 151; *anotinum*, 170; *clavatum*, 54, 169, 170, var. *megastachyon*, 54, 103, 116; *complanatum*, 116, 137, 169-171, 211, 212, near *Hartland*, Vermont, 211, var. *Chamaecyparissus*, 169, var. *flabelliforme*, 128, 139, 168; **flabelliforme**, 168-171; *inundatum*, 79, 128; *obscurem*, 169, 170, var. *den-droideum*, 54; *sabinaefolium*, 147; *Selago*, 136, 147; *tristachyum*, 169, 170.  
*Lycopus rubellus*, 8; *uniflorus*, 139.  
*Lygodium palmatum*, 78.  
*Lyonia calyculata*, 215.  
*Lysimachia terrestris*, 139.  
  
*Magnolia virginica*, 150, 155.  
 Maine, Winter Meeting of the Josselyn Botanical Society of, 90.  
 Maple, 57, 86; Swamp, 86.  
 Marigold, Marsh, 123.  
 Marsh Marigold, 123.  
*Marsilea*, 221; *quadrifolia*, 79, 221.  
 Maryland and Virginia, The Typhae of, 241; *Solidago polycephala* in, 106.  
 Massachusetts and New Jersey, A new Species of *Scirpus* from, 4; *Castanea pumila* in eastern, 90; *Cynanchum nigrum* in Barnstable, 70; *Dracocephalum thymiflorum* a casual Plant at Westford, 212; Pteridophyte Notes from eastern, 101; Some fleshy Fungi of Stow, 57; Two lost Carices from eastern, 243.  
 Massey, A. B., Plants in Flower February 8, 1911, at Clemson College, South Carolina, 91.  
*Matricaria Chamomilla*, 90; *inodora*, 90; *suaveolens*, 90.  
*Medeola*, 217.  
*Melampsora*, 14.  
  
*Melampyrum lineare*, 117, 149.  
*Melanopus*, 64.  
*Melilotus alba*, 89; *officinalis*, 89.  
*Microstylis monophyllos*, 117, 125; *unifolia*, 78.  
*Milium*, 173; *effusum*, 125.  
*Mimulus*, 18; *moschatus*, 115; *ringens*, 141.  
 Missouri *Rhexias*, 166.  
*Mitchella*, 216; *repens*, 32, 216, forma **leucocarpa**, 32.  
*Mitella nuda*, 125, 217.  
*Monotropa*, 17; *Hypopitys*, 18; *uniflora*, 18.  
*Montia lamprosperma*, 122; *rivularis*, 140.  
 Moore, A. H., *Solidago polycephala* in Maryland, 106.  
*Morchella esculenta*, 58.  
 Mosses, Notes on Connecticut,—II, 40.  
 Mountain Ash, 238; Cranberry, 136.  
*Myrica cerifera*, 243.  
*Myriophyllum humile*, 18; *spicatum*, 182; *tenellum*, 135, 140.  
  
*Najas guadalupensis*, 151, 160.  
*Nanomitrium Austini*, 46.  
 Nantucket, Three additional Plants from, 105.  
*Naucoria*, 57; *firma*, 64.  
 Necessary Change of Name, 55.  
*Nemopanthus mucronata*, 78.  
*Nepeta*, 18.  
 New England Botanical Club, Notes from the Phaenogamic Herbarium of the,—II, 177; Federation of Natural History, 187.  
 Newfoundland and southern Labrador, A botanical Expedition to, 109.  
 New Hampshire, *Helianthus subrhomboides* in, 103; A *Scirpus* new to, 55.  
 New Jersey, A new Species of *Scirpus* from Massachusetts and, 4.  
 New Species of *Scirpus* from Massachusetts and New Jersey, A, 4; Textbook on Botany, A, 16; Variety of *Rubus canadensis*, 193.  
 Nichols, G. E., Notes on Connecticut Mosses,—II, 40.  
 North America, Harshberger's Phytogeographic Survey of, 213.  
 Northern Variety of *Gaylussacia dumosa*, 95.  
 Norway Pine, 139.  
 Notes from eastern Massachusetts, Pteridophyte, 101; from my Herbarium, 71; from the Phaeno-



- gamie Herbarium of the New England Botanical Club, — II, 177; on algae, — X, 184; on Botrychium from Tennessee, 14; on certain Leguminosae, 33; on Connecticut Mosses, — II, 40; on Connecticut Plants, 30; on the Genus *Cirsium*, 238.
- Nymphaea*, 21; *advena*, 20.
- Oak, 57, 86.
- Observations on some Plants of eastern Connecticut, 68.
- Odontites rubra*, 182.
- Oenothera*, 209; *biennis*, 209; *grandiflora*, 210; *Oakesiana*, 218; *pumila*, 139; Systematic Studies on, — II, *Oenothera Tracyi*, sp. nov., 209; **Tracyi**, 210, 211.
- Omphalia pyxidata*, 58.
- On Gynodioecism in *Plantago lanceolata*, 199.
- Onagra*, 209.
- Onoclea sensibilis*, 126, 139.
- Ophioglossum vulgatum*, 79.
- Opuntia vulgaris*, 151.
- Orchids, 217.
- Orchis rotundifolia*, 115, 221.
- Ornithological observations on *Cleistogamy*, 76.
- Orontium aquaticum*, 151.
- Orthotrichum pumilum*, 46.
- Oryzopsis asperifolia*, 139.
- Osmorhiza*, 18; *divaricata*, 148; *obtusata*, 115, 126, 138.
- Osmunda*, 134; *cinnamomea*, 139; *regalis*, 132, var. *pumila*, 133.
- Osterhout, W. J. V., A new Textbook on Botany, 16.
- Ostrya virginica*, 141, 148.
- Oxybaphus nyctagineus*, 89.
- Oxytropis campestris*, 218, var. *caerulea*, 119, 129, 216.
- Panax quinquefolium*, 218.
- Panicum*, 66, 173–175; *adspersum*, 67; *albemarlense*, 67; *albomarginatum*, 68; *amaroides*, 67; *amarulum*, 67; *amarum*, 67; *auburne*, 67; *barbipulvinatum*, 67; *Bicknellii*, 67; *boreale*, 139; *Boscii*, 76; *bulbosum*, 9, 10; *caerulescens*, 67; *calliphyllum*, 67; *capillare*, 173, *clandestinum*, 76; *Clutei*, 67; compared with that in Grays' Manual, Recent Treatment of, 66; *cryptanthum*, 68; *dichotoma*, 176; *dichotomiflorum*, 175; *dichotomum*, 173, 176; *Helleri*, 68; *Helopus*, 174; *hemitomum*, 68; *hemitomum*, 68; *hians*, 67; *implicatum*, 116, 135; *italicum*, 173; *Joorii*, 68; *languidum*, 68; *mala-cophyllum*, 68; *mattamuskeetense*, 67; *miliaceum*, 173–175; *miliare*, 175; *nitidum*, 67; *obtusum*, 68; *occidentale*, 175; *oricola*, 67; *ovale*, 68; *philadelphicum*, 54; *proliferum*, 175; *pseudopubescens*, 68; *pubescens*, 175; *rhizomatum*, 67; *roanokense*, 67; *scoparium*, 151, 175; *sphaerocarpon*, 68, var. *inflatum*, 68; *tenuis*, 66, 68; Type of the Genus, 173; *verrucosum*, 151; *versicolor*, 175; *violaceum*, 175; *virgatum*, 88, 175, var. *cubense*, 67, var. *obtusum*, 67; *Wrightianum*, 67.
- Paper Birch, 207.
- Paractaenium*, 174.
- Parnassia Kotzebuei*, 124; *parviflora*, 119, 138.
- Partridge-berry, 123, 216.
- Paspalum psammophilum*, 22.
- Pease, A. S., *Helianthus subrhomboides* in New Hampshire, 103.
- Pedicularis Furbishiae*, 218.
- Peltandra*, 179; *virginica*, 179.
- Penhallow, David Pearce, [Notice of], 1; A correction regarding Professor, 56.
- Penicillium glaucum*, 86.
- Pennisetum*, 175.
- Penthorum sedoides*, 141.
- Petasites palmata*, 125.
- Phanopyrum*, 174.
- Phegopteris Dryopteris*, 124, 125, 131; *polypodioides*, 125; *Robertiana*, 131, 219.
- Philonotis*, 45; *caespitosa*, 46, var. *compacta*, 46, var. *heterophylla*, 46, var. *laxa*, 46.
- Phleum alpinum*, 136; *pratense*, 208.
- Phryma leptostachya*, 218.
- Phyllodoce caerulea*, 133, 136.
- Physalis grandiflora*, 219; *pruinosa*, 19.
- Phytogeographic Survey of North America, Harshberger's, 213.
- Picea alba*, 217, 219; *canadensis*, 137, 217, 219; *mariana*, 54, 138.
- Pine, 57; Norway, 139; Red, 139, 161.
- Pinguicula vulgaris*, 115, 124, 136.
- Pinus Banksiana*, 217, 219; *divaricata*, 217; *Parryana*, 229; *resinosa*, 116, 139, 146, 161; *rigida*,

- 103, 253; *Strobilus*, 77, 139, 219, 253, stump-healing in, 253.
- Plantago aristata*, 19, var. *Nuttallii*, 19; *decipiens*, 216; *lanceolata*, 199–203, 205, 206, var. *alopecuroides*, 204, On *Gynodioecism* in, 199, var. *sphaerostachya*, 204; *maritima*, 215; *media*, 200.
- Plants from Nantucket, Three additional, 105; in Flower February 8, 1911, at Clemson College, South Carolina, 91; Notes on Connecticut, 30; of Connecticut, Some introduced, 88; of eastern Connecticut, Observations on some, 68.
- Plumbago, 123.
- Pluteolus callistus*, 65.
- Poa alpina*, 119, 136; *eminens*, 119, 123; *Sandbergi*, 141.
- Podostemum ceratophyllum*, 221.
- Pogonia ophioglossoides*, 139, 220.
- Poisonous Dogwood, 57; Ivy, 57.
- Polygala Nuttallii*, 8.
- Polygonaceae, 89.
- Polygonella articulata*, 180.
- Polygonum cilinode*, 141, 148; *Muhlenbergii*, 18; *ramosissimum*, 89; *Roberti*, 138; *sagittatum*, 18, 139; *Fowleri*, 180.
- Polyides*, 187; *rotundus*, 187.
- Polypodium vulgare*, 105.
- Polyporaceae, 64.
- Polyporus*, 63; *admirabilis*, 63–65; *squamosus*, 64; *varius*, 64.
- Polystichum Braunii*, 116, 218, 238; *Lonchitis*, 116, 117.
- Polytrichum*, 91; *alpinum*, 46; *strictum*, 46.
- Poplar, Black, 195; Virginian Black, 198.
- Populus*, 197; *alba*, var. *Bolleana*, 197; *angulata*, 12, 13; *balsamifera*, 138, 147; *canadensis*, 13, 196, 199; *candicans*, 237; *deltoides*, 12–14, 195–197, 199; *grandidentata*, 141, 148; *marilandica*, 196, 199; *monilifera*, 12, 197, 198; *nigra*, var. *italica*, 197; var. *virginiana*, 198; *tremuloides*, 117; *virginiana*, 12–14, 195–198, and *P. angulata*, 12.
- Potamogeton crispus*, 221; *filiformis*, 138; *Oakesianus*, 134, 135, 140, 148, 160; *pectinatus*, 177; *praelongus*, forma *elegans*, 117; *Vaseyi*, 71.
- Potentilla Anserina*, 180; *canadensis*, var. *simplex*, 106; *fruticosa*, var. *tenuifolia*, 129; *intermedia*, 89; *maculata*, 127, 137; *nivea*, 127; *recta*, 89; *tridentata*, 238.
- Prenanthes altissima*, 141, 148; *mainensis*, 218; *racemosa*, 148.
- Primula farinosa*, var. *macropoda*, 138, 216.
- Pringle, Cyrus Guernsey, 225.
- Proserpinaca pectinata*, 8.
- Prunus americana*, var. *mollis*, 31; *maritima*, 91; *nigra*, 31; *Persica*, 91; *pumila*, 141.
- Psalliotia*, 62.
- Ptelea*, 81; *mollis*, 80–82, var. **cryptoneura**, 81, 82, a Wafer-ash of the Georgia Sand-hills, 80; *trifoliata*, 80, var. *mollis*, 80, 81.
- Pteridophyte Notes from eastern Massachusetts, 101.
- Pterigynandrum filiforme*, 46.
- Pubescent Variety of *Vaccinium vacillans*, 235.
- Puccinellia*, 216.
- Pycnanthemum Torrei*, 32.
- Pyrola americana*, 97, 139, 146; *asarifolia*, 115, 141, 148, var. *incarnata*, 138, 141; *elliptica*, 147; *minor*, 138; *secunda*, 126; *uliginosa*, 221.
- Pyrus americana*, 238; *arbutifolia*, var. *atropurpurea*, 117, 135, 149; *melanocarpa*, 139; *sitchensis*, 138, 238.
- Quercus*, 229; *alba*, 217, 219; *coccinea*, 219; *prinoides*, 219; *rubra*, 141; *velutina*, 217.
- Racomitrium aciculare*, 43; *fasciculare*, 43.
- Radicula aquatica*, 229.
- Range of the Black Birch to be restricted, 206.
- Ranunculus abortivus*, 125; *affinis*, 223; *Allenii*, 146, 223; *Harveyi*, 146; *hyperboreus*, 122; *Macounii*, 138; *pedatifidus*, 223; *pennsylvanicus*, 180; *repens*, 19; *sceleratus*, 19.
- Recent Treatment of *Panicum* compared with that of Gray's Manual, 66.
- Red Berry, 123; Pine, 139, 161.
- Reports on the Flora of the Boston District,—VIII, 27, IX, 72, X, 82, XI, 104, XII, 232, XIII, 248.
- Reynolds, E. S., Notes on *Botrychium* from Tennessee, 14.

- Rhexia, 166; **latifolia**, 167; *mariana*, 166-168; *virginica*, 167, 168.  
 Rhexias, The Missouri, 166.  
 Rhinanthus, 116.  
 Rhododendron, 86; *canadense*, 79; *lapponicum*, 136; *Rhodora*, 220, 221.  
 Rhodora, 221.  
 Rhus Toxicodendron, 57; *Vernix*, 57.  
 Rhytidadelphus squarrosus, 46.  
 Ribes hirtellum, 73-76, var. **callicola**, 76, var. **saxosum**, 76, 138, The Varieties of, 73; *inermis*, 74-76; *lacustre*, 138; *oxyacanthoides*, 74, 75, var. **callicola**, 74-76, var. **saxosum**, 75, 76; *prostratum*, 138, 216; **saxosum**, 75, 76; *triste*, 125, 138.  
 Riddle, L. W., An important Publication on Lichens, 15.  
 Robinson, B. L., *Erucastrum Pollichii* adventive in America, 10; Notes on the Genus *Cirsium*, 238.  
 Rosa acicularis, var. *Bourgeauiana*, 180; *canina*, 31; *nitida*, 31; *rugosa*, 71; *virginiana*, 139, 146, 149.  
 Rosaceae, 106, 229.  
 Rotala ramosior, 8.  
 Rubus, 216; *amabilis*, 56; **amicalis**, 56; *arcticus*, 123; *canadensis*, 55, 194, A new Variety of, 193, var. **septemfoliolatus**, 193; *Chamaemorus*, 95, 99, 114, 123; *recurvus*, 194.  
 Rudbeckia, 18.  
 Rumex altissimus, 89; *hastatus*, 151; *mexicanus*, 89; *occidentalis*, 123; *persicarioides*, 179.  
 Russula, 57, 62; *abietina*, 62, 65; *obscura*, 59, 60; *palustris*, 59, 60, 65.  
 Rynchospora, 104; *alba*, 104, var. *macra*, 104; *capillacea*, 30; *fusca*, 105, 136, 139; *glomerata*, 105; *macrostachya*, 105, var. *inundata*, 105; *Torreyana*, 151.  
 Sabatia chloroides, 150; *dodecandra*, 150, 159, 161.  
 Sagina decumbens, 8, 151; *occidentalis*, 219.  
 Sagittaria arifolia, 141; *Engelmanniana*, 247; *latifolia*, 18, 141.  
 Salicaceae, 196.  
 Salix, 147; *balsamifera*, 138; *Barclayi*, 138; **callicola**, 251, 252, 253, a little known northern Willow, 251; *candida*, 114, 123, 138; *cordifolia*, 123; *discolor*, 139; *glaucophylla*, 138; *humilis*, 117, 139, 146; *lanata*, 127, 252, 253, var. *Macouniana*, 251, 252; *lucida*, var. *intonsa*, 116; *Macounii*, 252; *pedicellaris*, var. *hypoglaucha*, 116; *pellita*, 179; *petiolaris*, 111; *Pseudomyrsinites*, 119, 123; *reticulata*, 110, 127; *Richardsoni*, 252, var. *Macouniana*, 251, 252; *Uva-ursi*, 123, 136; *vestita*, 117, 123, 129.  
 Salsola Kali, var. *tenuifolia*, 89.  
 Sambucus canadensis, 141.  
 Samolus floribundus, 182.  
 Sanicula, 18; *gregaria*, 218; *marilandica*, 149.  
 Sanders, A. L., A striking color Form of *Viola pedata*, 172.  
 Sargassum, 187; *vulgare*, 187.  
 Sarracenia, 114, 221; *purpurea*, 78, 220.  
 Saururus cernuus, 151.  
 Saxifraga, 18; *aizoides*, 117; *Aizoon*, 117, 238; *caespitosa*, 119, 216; *Geum*, 140; *oppositifolia*, 117.  
 Scheuchzeria palustris, 116.  
 Schizaea filifolia, 110; *pusilla*, 110, 133-135, 140, 148, 159, 161.  
 Scirpus, 4, 5, 7, 8, 21, 82, 83, 133; *americanus*, 83; *atrocinetus*, 4-7, 83, 138, var. *brachypodus*, 83, var. *grandis*, 7; *atrovirens*, 83; *caespitosus*, 95, 133, 136, 160; *campestris*, var. *Fernaldi*, 83, var. *novaeangliae*, 83, 178, var. *paludosus*, 83; *cyperinus*, 7, 21, 83, var. *Andrewsii*, 84, var. *condensatus*, 84, var. *pelius*, 84; *debilis*, 84, var. *Williamsii*, 84; *Eriophorum*, 7; *fluviatilis*, 84, 178; from Massachusetts and New Jersey, A new Species of, 4; *georgianus*, 84, 139; *Hallii*, 84, 151, 155; *heterochaetus*, 84; *hudsonianus*, 84, 138; **Longii**, 6-8, 84, 151; *nanus*, 84; new to New Hampshire, 55; *occidentalis*, 85; *Olneyi*, 85; *pauciflorus*, 138, 221; *pedicellatus*, 7, 85, var. *pullus*, 85; *planifolius*, 85; *robustus*, 85; *rubrotinctus*, 85, var. *confertus*, 55; *Smithii*, 30, 85, var. *setosus*, 30, 178; *subterminalis*, 85, 128, 135; *sylvaticus*, 85; *Torreyi*, 85; *validus*, 21, 85.  
 Scleria, 105; *reticularis*, 105, 151; *triglomerata*, 105; *verticillata*, 30.  
 Sclerolepis uniflora, 152.



- Scrophularia*, 115; *leporella*, 218; *nodosa*, 115.  
 Scudder, H., *Similarity of Color in Bud and Leaf*, 86.  
*Scurvy Grass*, 119, 124.  
*Scutellaria galericulata*, 146.  
*Sedum roseum*, 216, 237.  
*Selaginella selaginoides*, 115.  
*Senecio*, 116, 131; *discoideus*, 115; *pauciflorus*, 124; *Pseudo-Arnica*, 123; *Robbinsii*, 183, 221; *sylvaticus*, 183.  
*Setaria*, 174-176; *italica*, 176; *verticillata*, 88.  
*Shepherdia canadensis*, 138.  
*Sideritis montana*, 89.  
*Silene latifolia*, 21.  
 Silver Birch, 207.  
*Similarity of Color in Bud and Leaf*, 86.  
*Sisymbrium*, 11; *altissimum*, 10; *Erucastrum*, 11; *gallicum*, 11; *hirtum*, 11; *Loeselii*, 89; *Sophia*, 89.  
*Sium cicutae-folium*, 146.  
 Sixteenth annual Meeting of the Vermont Botanical Club, 237.  
 Skunk Currant, 138.  
*Smilacina stellata*, 216; *trifolia*, 78.  
*Smilax rotundifolia*, 179.  
 Societies, New England Federation of Natural History, 187.  
 Society of Maine, Winter Meeting of the Josselyn Botanical, 90.  
*Solanum carolinense*, 89.  
*Solidago*, 19, 106; *bicolor*, 19; *canadensis*, 32; *Cutleri*, 238; *Elliottii*, 78; *floribunda*, 107; *graminifolia*, var. *Nuttallii*, 107; *junceae*, 148; *latifolia*, 148; *macrophylla*, 125, 138, var. *thyrsoides* 71; *multiradiata*, 119; *polyccephala*, 106, 107, in Maryland, 106; *puberula*, 19; *racemosa*, 183; *Randii*, 238; *resinosa*, 146; *rugosa*, 139, 183, var. *villosa*, 131; *squarrosa*, 141, 148, 161; *ulmifolia*, 183; *unguiculata*, 135.  
*Solidago* § *Euthamia*, 106.  
 Some boreal Species and Varieties of *Antennaria* and *Anaphalis*, 23; *fleshy Fungi* of Stow, Massachusetts, 57; *introduced Plants of Connecticut*, 88.  
 South Carolina, *Plants in Flower* February 8, 1911, at Clemson College, 91.  
*Sparganium diversifolium*, 148; *hyperboreum*, 134, 160.  
*Spartina*, 21; *cynosuroides*, 21, 22; *glabra*, var. *alterniflora*, 128.  
 Species and Varieties of *Antennaria* and *Anaphalis*, Some boreal, 23; of *Scirpus* from Massachusetts and New Jersey, A new, 4.  
 Speedwell, 124.  
*Spergularia rubra*, 135, 140.  
*Sphagnum*, 96, 220, 221.  
*Spiraea*, 18; *latifolia*, 18, 139, 146.  
*Spiranthes Beckii*, 8; *Romanzoffiana*, 138.  
 Spruce, 57; Black, 114; White, 114.  
*Stachys ambigua*, 151.  
*Statice*, 134, 216; *sibirica*, 118.  
*Steinchisma*, 174; *hians*, 67.  
*Steironema ciliatum*, 141, 148.  
*Stellaria borealis*, 78; *crassifolia*, 122; *media*, 91; *uliginosa*, 237.  
*Stenophyllus*, 73; *capillaris*, 73.  
*Stipa canadensis*, 133.  
 Stow, Massachusetts, Some fleshy Fungi of, 57.  
 Strand Wheat, 122, 123.  
*Streptopus amplexifolius*, 125, 237; *longipes*, 237, An Addition to the Description of, 237; *roseus*, 237.  
 Striking color Form of *Viola pedata*, 172.  
 Stump-healing in *Pinus Strobus*, 253.  
*Suaeda*, 18; *linearis*, 18, 180; *maritima*, 18.  
 Subterranean Organs of *Cinna arundinacea*, 9.  
 Swamp Maple, 86.  
 Sweet Birch, 206.  
 Systematic Studies on *Oenothera*,—I. *Oenothera Tracyi*, sp. nov., 209.  
 Tamarack, 57.  
*Tanacetum huronense*, 127, 138, 218.  
*Taraxacum ceratophorum*, 117; *erythrospermum*, 18.  
*Taxus canadensis*, 215.  
 Tennessee, Notes on *Botrychium* from, 14.  
 Teratology in *Trillium ovatum* Pursh, 189.  
*Thalictrum*, 18; *alpinum*, 117, 129, 136, 223; *confine*, 147; *dioicum*, 111, 146, 147, 218; *divicium*, 146; *polygamum*, 147; *purpurascens*, 146, 147.  
 Thistle, 116.  
*Thlaspi arvense*, 89.  
 Thompson, E. J., *Botanizing in central Connecticut*, 77.

- Three additional Plants from Nantucket, 105.
- Thuja, 221; *occidentalis*, 115, 142, 161, 237.
- Tiarella, 193; *cordifolia*, 141, 192, Dimorphism in, 192.
- Tidestrom, I., *Populus virginiana* Fouger, 195; The Typhae of Maryland and Virginia, 241.
- Tilia cordata, 181.
- Tithymalopsis, 164.
- Tithymalus, 164.
- Toffieldia glutinosa, 115, 138; *palustris*, 117, 129, 136.
- Townsend, C. W., A Comment on the Use of the Term Labrador in Natural History, 236.
- Tricholoma, 63; *ustale*, 63, 65, 66.
- Trifolium dubium, 34; *procumbens*, 34.
- Triglochin palustris, 138.
- Trillium, 189-191; *cernuum*, 139; *erectum*, 141; *grandiflorum*, 191; *ovatum*, 189, 191, Teratology in, 189; *recurvatum*, 191, 216; *sessile*, 191; *undulatum*, 141.
- Triosteum aurantiacum, 218.
- Trisetum melicoides, 132, 138; *spicatum*, 177.
- Trollius laxus, 31.
- Turritis laevigatus, 38.
- Two lost Carices of eastern Massachusetts, 243.
- Type of the Genus Panicum, 173.
- Typha, 242; *angustifolia*, 241-243, var. **virginica**, 242; *glauca*, 243; *latifolia*, 241-243.
- Typhae of Maryland and Virginia, 241.
- Urochloa, 174; *panicoides*, 174; *uniseta*, 175.
- Urtica, 18, 123.
- Usnea trichodea, 91.
- Utricularia clandestina, 78, 116, 135, 140; *cornuta*, 220; *resupinata*, 247.
- Vaccinium, 216, 235; *caespitosum*, 218; *corymbosum*, 54; *dumosum*, 96-98; *frondosum*, 96, 99; *hirtellum*, 96, 97; *macrocarpon*, 54, 117, 135, 139, 149, 217, 220; *ovalifolium*, 116; *Oxycoccus*, 220; *pennsylvanicum*, var. *angustifolium*, 96, var. *myrtilloides*, 235; *uliginosum*, 123; *vacillans*, 235, var. **crinitum**, 235, A pubescent Variety of, 235; *Vitis-Idaea*, var. *minus*, 123, 136, 182, 216.
- Valeriana septentrionalis, 221; *sylvatica*, 116; *uliginosa*, 115, 116.
- Vallisneria, 215; *spiralis*, 111, 215.
- Variations of Lathyrus palustris in eastern America, 47.
- Varieties of Antennaria and Anaphalis, Some boreal Species and, 23; of Ribes hirtellum, 73.
- Variety of Fragaria virginiana, A boreal, 106; of Gaylussacia dumosa, The northern, 95; of Rubus canadensis, A new, 193; of Vaccinium vacillans, A pubescent, 235.
- Veratrum viride, 141, 148.
- Verbena stricta, 89.
- Vermont Botanical Club, Sixteenth annual Meeting of the, 237; Botanical Club, The winter Meeting of the, 87; Glyceria acutiflora in, 92; Lycopodium complanatum near Hartland, 211.
- Veronica, 18; *Anagallis-aquatica*, 32; *humifusa*, 124.
- Vesicaria arctica, 223.
- Vetch, Kidney, 240.
- Viburnum acerifolium, 111, 216; *alnifolium*, 141, 161; *cassinoides*, 111, 139, 146, 216; *Lentago*, 111, 216; *pauciflorum*, 138, 216; *venosum*, 151.
- Vicia, 18; *Cracca*, 34, 89; *hirsuta*, 34; *tetrasperma*, 33; *villosa*, 34.
- Viola, 110, 182; *adunca*, 147; *arenaria*, 147; *Brittoniana*, 182; *cucullata*, 149; *labradorica*, 147; *nephrophylla*, 138; *pedata*, 172, *forma alba*, 172, var. *lineariloba*, 172, *forma rosea*, 172, A striking color Form of, 172; *primulaefolia*, 218; *renifolia*, 125; *rostrata*, 146, 147; *Selkirkii*, 125, 126; *septentrionalis*, 141.
- Virginia, The Typhae of Maryland and, 241.
- Virginian Black Poplar, 198.
- Wafer-ash, 80.
- Waldsteinia fragarioides, 218.
- Ware, R. A., Another Station for Centaurea diffusa, 100.
- Water Lily, 191.
- Westford, Massachusetts, Dracopcephalum thymiflorum a casual Plant at, 212.
- Wheat, Strand, 122, 123.
- White Spruce, 114.

- Whortle-berry, 98, 123.
- Wiegand, K. M., A boreal Variety of *Fragaria virginiana*, 106; *Cornus canadensis*, var. *intermedia* in eastern America, 107; *Epilobium palustre*, var. *longirameum*, 188; *Salix calcicola*, a little known northern Willow, 251; Some boreal species and Varieties of *Antennaria* and *Anaphalis*, 23.
- Williams, E. F., *Castanea pumila* in eastern Massachusetts, 90; Notes from my Herbarium, 71.
- Willow, 86; *Salix calcicola*, a little known northern, 251.
- Winter Meeting of the Josselyn Botanical Society of Maine, 90; Meeting of the Vermont Botanical Club, 87.
- Woodsia glabella*, 129, 218, 238; *hyperborea*, 218, 219; *obtusa*, 101; *scopulina*, 219.
- Woodward, R. W., Observations on some Plants of eastern Connecticut, 68.
- Xanthium spinosum*, 89.
- Xyris caroliniana*, 54; *montana*, 159; *Smalliana*, 247.
- Yellow Birch, 206, 207; Rocket, 123.
- Zigadenus chloranthus*, 138.
- Zostera marina*, 111, 215.



